PCT

'ORLD INTELLECTUAL PROPERTY ORGANIZAT. International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:

C12N 15/29, C07K 14/415, C12N 5/04, 5/14, A01H 5/00, C12N 15/10, 15/82

(11) International Publication Number: WO 99/19492

(43) International Publication Date: 22 April 1999 (22.04.99)

(21) International Application Number: PCT/EP98/06977

(22) International Filing Date: 9 October 1998 (09.10.98)

(30) Priority Data:

PO 9745 10 October 1997 (10.10.97) AU

(71) Applicant (for all designated States except US):
RHONE-POULENC AGRO [FR/FR]; 14/20, rue Pierre
Baizet, F-69009 Lyon (FR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): DOUTRIAUX, Marie-Pascale [FR/FR]; 64, route de Villebon, F-91160 Saulx les Chartreux (FR). BETZNER, Andreas, Stefan [AU/AU]; 40 Dallachy Place, Page, ACT 2614 (AU), FREYSSINET, Georges [FR/FR]; 21, rue de Nervieux, F-69450 Saint Cyr au Mont d'Or (FR). PEREZ, Pascal [FR/FR]; 17, chemin de la Pradelle, Varennes, F-63450 Chanonat (FR).

(74) Agent: GENIN, Patrick; Rhône-Poulenc Agro, DPI, 14/20, rue Pierre Baizet, F-69009 Lyon (FR). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN,

Published

TD, TG).


Without international search report and to be republished upon receipt of that report.

(54) Title: METHODS FOR OBTAINING PLANT VARIETIES

(57) Abstract

An isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide functionally involved in the DNA mismatch repair system of a plant.






FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenía
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senega!
ΑU	Australia	GA	Gabon	LV	Latvia	S2	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajíkistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
ВG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
ВJ	Benin	ΙE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Сапада	IT	Italy	MX	Mexico	υz	Uzbekistan
CF	Central African Republic	JР	Japan	NE	Niger	VN	Viet Nam
ÇG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
cυ	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:

C12N 15/29, C07K 14/415, C12N 5/04,
5/14, A01H 5/00, C12N 15/10, 15/82

(11) International Publication Number: WO 99/19492

(43) International Publication Date: 22 April 1999 (22.04.99)

(21) International Application Number: PCT/EP98/06977

(22) International Filing Date: 9 October 1998 (09.10.98)

(31) International Publication Date: 22 April 1999 (22.04.99)

(43) International Publication Date: 22 April 1999 (22.04.99)

(51) International Publication Number: PCT/EP98/06977

(51) International Publication Number: WO 99/19492

(52) International Publication Number: 22 April 1999 (22.04.99)

(53) International Publication Number: 32 April 1999 (22.04.99)

(54) International Publication Number: 32 April 1999 (22.04.99)

(55) International Publication Number: 32 April 1999 (22.04.99)

ΑU

(71) Applicant (for all designated States except US):
RHONE-POULENC AGRO [FR/FR]; 14/20, rue Pierre
Baizet, F-69009 Lyon (FR).

10 October 1997 (10.10.97)

(72) Inventors; and

(30) Priority Data:

PO 9745

(75) Inventors/Applicants (for US only): DOUTRIAUX, Marie-Pascale [FR/FR]; 64, route de Villebon, F-91160 Saulx les Chartreux (FR). BETZNER, Andreas, Stefan [AU/AU]; 40 Dallachy Place, Page, ACT 2614 (AU). FREYSSINET, Georges [FR/FR]; 21, rue de Nervieux, F-69450 Saint Cyr au Mont d'Or (FR). PEREZ, Pascal [FR/FR]; 17, chemin de la Pradelle, Varennes, F-63450 Chanonat (FR).

(74) Agent: GENIN, Patrick; Rhône-Poulenc Agro, DPI, 14/20, rue Pierre Baizet, F-69009 Lyon (FR).

BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

Without international search report and to be republished upon receipt of that report.

(54) Title: METHODS FOR OBTAINING PLANT VARIETIES

(57) Abstract

An isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide functionally involved in the DNA mismatch repair system of a plant.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

	AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
	AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
	AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
	ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
	AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
	BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
	BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
	BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
	BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
	BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
	ВJ	Benin	1E	Ireland	MN	Mongolia	UA	Ukraine
	BR	Brazil	IL.	Israel	MR	Mauritania	UG	Uganda
	BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
	CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
	CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
	CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
	CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
i	CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
	CM	Cameroon		Republic of Korea	PL	Poland		
	CN	China	KR	Republic of Korea	PT	Portugal		
	CU	Cuba	КZ	Kazakstan	RO	Romania		
l	CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
l	DE	Germany	LI	Liechtenstein	SD	Sudan		
l	DK	Denmark	LK	Sri Lanka	SE	Sweden		
l	EE	Estonia	LR	Liberia	SG	Singapore		

WO 99/19492

10

1

Methods for Obtaining Plant Varieties

TECHNICAL FIELD

The present invention relates to nucleotide sequences which encode polypeptides involved in the DNA mismatch repair systems of plants, and to the polypeptides encoded by those nucleotide sequences. The invention also relates to nucleotide sequences and polypeptide sequences for use in altering the DNA mismatch repair system in plants. The invention also relates to a process for altering the DNA mismatch repair system of a plant cell, to a process for increasing genetic variations in plants and to processes for obtaining plants having a desired characteristic.

BACKGROUND OF THE INVENTION

Plant breeding essentially relies on and makes use of genetic variation which occurs naturally within and between members of a family, a genus, a species or a subspecies. Another source of genetic variation is the introduction of genes from other organisms which may or may not be related to the host plant.

Allelic loci or non-allelic genes which constitute or contribute to desired quantitative (e.g. growth performance, yield, etc.) or qualitative (e.g. deposition, content and composition of seed storage products; pathogen resistance genes: etc.) traits that are absent, incomplete or inefficient in a species or subspecies of interest are typically introduced by the plant breeder from other species or subspecies, or *de novo*. This introduction is often done by crossing, provided that the species to be crossed are sexually compatible. Other means of introducing genomes, individual chromosomes or genes into plant cells or plants are well known in the art. They include cell fusion, chemically aided transfection (Schocher et al., 1986, Biotechnology 4: 1093) and ballistic (McCabe et al., 1988, Biotechnology 6: 923), microinjection (Neuhaus et al., 1987, TAG 75: 30), electroporation of protoplasts (Chupeau et al., 1989, Biotechnology 7: 53) or microbial transformation methods such as Agrobacterium mediated transformation (Horsch et al., 1985, Science 227: 1229; Hiei et al., 1996, Biotechnology 14: 745).

However, when a foreign genome, chromosome or gene is introduced into a plant, it will often segregate in subsequent generations from the genome of the recipient plant or plant cell during mitotic and meiotic cell divisions and, in consequence, become lost from the host plant or plant cell into which it had been introduced. Occasionally, however, the introduced genome, chromosome or gene physically combines entirely or in part with the genome, chromosome or gene of the host plant or plant cell in a process which is called recombination.

Recombination involves the exchange of covalent linkages between DNA molecules in regions of identical or similar sequence. It is referred to here as homologous recombination if donor and recipient DNA are identical or nearly identical (at least 99%)

hase sequence identity), and as homeologous recombination if donor and recipient DNA are not identical but are similar (less than 99% base sequence identity).

The ability of two genomes, chromosomes or genes to recombine is known to depend largely on the evolutionary relation between them and thus on the degree of sequence similarity between the two DNA molecules. Whereas homologous recombination is frequently observed during mitosis and meiosis, homeologous recombination is rarely or never seen.

From a breeder's perspective, the limits within which homologous recombination occurs, therefore, define a genetic barrier between species, varieties or lines, in contrast to homeologous recombination which can break this barrier. Homeologous recombination is thus of great importance for plant breeding. Accordingly there is a need for a process for enhancing the frequency of homeologous recombination in plants. In particular, there is a need for a process of increasing homeologous recombination to significantly shorten the length of breeding programs by reducing the number of crosses required to obtain an otherwise rare recombination event.

At least in Escherichia coli. homologous and homeologous recombination are known to share a common pathway that requires among others the proteins RecA, RecB, RecC, RecD and makes use of the SOS induced RuvA and RuvB, respectively. It has been suggested that mating induced recombination follows the Double-Strand Break Repair 20 model (Szostak et al., 1983, Cell 33, 25-35), which is widely used to describe genetic recombination in eukaryotes. Following the alignment of homologous or homeologous DNA double helices the RecA protein mediates an exchange of a single DNA strand from the donor helix to the aligned recipient DNA helix. The incoming strand screens the recipient helix for sequence complementarity, seeking to form a heteroduplex by hydrogen 25 bonding the complementary strand. The displaced homologous or homeologous strand of the recipient helix is guided into the donor helix where it base pairs with its counterpart strand to form a second heteroduplex. The resulting branch point then migrates along the aligned chromosomes thereby elongating and thus stabilising the initial heteroduplexes. Single stranded gaps (if present) are closed by DNA synthesis. The strand cross overs 30 (Holliday junction) are eventually resolved enzymatically to yield the recombination products.

Although in wild type E. coli homologous and homeologous recombination are thus mechanistically similar if not identical, homologous recombination in conjugational crosses E. coli x E. coli occurs five orders of magnitude more frequently than homeologous recombination in conjugational crosses E. coli x S. typhimurium (Matic et al. 1995; Cell 80, 507-515). The imbalance in favour of homologous recombination was shown to be caused largely by the bacterial MisMatch Repair (MMR) system since its

inactivation increased the frequency of homeologous recombination in *E. coli* up to 1000 fold (Rayssiguier et al. 1989, Nature 342, 396-401).

In E. coli, the MMR system (reviewed by Modrich 1991, Annual Rev Genetics 25, 229-253) is composed of only three proteins known as MutS, MutL and MutH. MutS recognizes and binds to base pair mismatches. MutL then forms a stable complex with mismatch bound MutS. This protein complex now activates the MutH intrinsic single stranded endonuclease which nicks the strand containing the misplaced base and thereby prepares the template for DNA repair enzymes.

During recombination, MMR components inhibit homeologous recombination. In vitro experiments demonstrated that MutS in complex with MutL binds to mismatches at the recombination branch point and physically blocks RecA mediated strand exchange and heteroduplex formation (Worth et al., 1994; PNAS 91, 3238-3241). Interestingly, the SOS dependent RuvAB mediated branch migration is insensitive to MutS/MutL, explaining the observed slight increase in SOS dependent homeologous recombination.

15 Homeologous mating even induces the SOS response, thereby taking advantage of RuvAB induction (Matic et al. 1995, Cell 80, 507-515).

The MMR system thus appears to be a genetic guardian over genome stability in *E. coli*. In this role it essentially determines the extent to which genetic isolation, that is, speciation, occurs. The diminished sensitivity of the SOS system to MMR, however, allows (within limits) for rapid genomic changes at times of stress, providing the means for fast adaptation to altered environmental conditions and thus contributing to intraspecies genetic variation and species evolution.

The important role of MMR in preserving genomic integrity has been established also in certain eukaryotes. In its efficiency, the human MMR, for example, may even counteract potential gene therapy tools such as triple-helix forming oligonucleotides including RNA-DNA hybrid molecules (Havre et al., 1993, J. Virology 67: 7234-7331; Wang et al., 1995, Mol. Cell. Biol. 15: 1759-1768; Kotani et al., 1996, Mol. Gen. Genetics 250: 626-634; Cole-Strauss et al., 1996, Science 273: 1387-1389). Such oligonucleotides are designed to introduce single base changes into selected DNA target sequences in order to inactivate for example cancer genes or to restore their normal function. The resulting base mismatches however are recognised by the mismatch repair system which then directs removal of the mismatched base, thereby reducing the efficiency of oligonucleotide induced site-specific mutagenesis.

To date, two families of related genes, homologous to the bacterial MutS and MutL genes have been identified or isolated in yeast and mammals (recent reviews by Arnheim and Shibata, 1997, Curr. Opinion Genet. Dev. 7, 364-370; Modrich and Lahue, 1996, Annual Rev. Biochem. 65, 101-133; Umar and Kunkel, 1996, Eur. J. Biochem. 238, 297-307). Biochemical and genetic analysis indicated that eukaryotic MutS homologs (MSH)

and MutL homologs (MLH, PMS), respectively, fulfil similar protein functions as their bacterial counterparts. Their relative abundance, however, could reflect different mismatch specificity and/or specialisation for different tissues or organelles or developmental processes such as mitotic versus meiotic recombination.

To date, six different genes homologous to *MutS* have been isolated in yeast (yMSH), and their homologs have been found in mouse (mMSH) and human (hMSH), respectively. Encoded proteins yMSH2, yMSH3 and yMSH6 appear to be the main MutS homologs involved in MMR during mitosis and meiosis in yeast, where the complementary proteins MSH3 and MSH6 alternatively associate with MSH2 to recognise different mismatch substrates (Masischky et al., 1996, Genes Dev. 10, 407-420). Similar protein interactions have been demonstrated for the human homologs hMSH2, hMSH3 and hMSH6 (Acharya et al., 1996, PNAS 93, 13629-13634).

MutL homologs (MLH and PMS), recently reviewed by Modrich and Lahue (1996, Annual Rev. Biochem. 65, 101-133) have so far been found in yeast (yMLH1 and yPMS1), mouse (mPMS2) and human (hMLH1, hPMS1 and hPMS2). The hPMS2 is a member of a family of at least 7 genes (Horii et al., 1994, Biochem. Biophys. Res. Commun. 204, 1257-1264) and its gene product is most closely related to yPMS1. Prolla et al. (1994, Science 265, 1091-1093) presented evidence for yPMS1 and yMLH1 to physically associate with each other and, together, to interact with the MutS homolog yMSH2 to form a ternary complex involved in mismatch substrate binding.

However, while medical interest in mismatch repair has prompted extensive research on MMR in bacteria, yeast and mammals, MMR genes have not been isolated from higher plants prior to the present invention and no attempts to adjust the plant MMR to plant breeding needs have been reported.

SUMMARY OF THE INVENTION

25

According to a first embodiment of the invention, there is provided an isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide functionally involved in the DNA mismatch repair system of a plant. In one form of this embodiment, the invention provides an isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human. More particularly, the invention provides polynucleotide sequences encoding polypeptides which are homologous to the mismatch repair polypeptides MSH3 and MSH6 of Saccharomyces cerevisiae. Still more particularly, the invention provides the coding sequences of the genes AtMSH3 and AtMSH6 of Arabidopsis thaliana, as defined hereinbelow, and polynucleotide sequences encoding polypeptides which are homologous to polypeptides encoded by AtMSH3 and AtMSH6.

According to a second embodiment of the invention, there is provided an isolated and purified polypeptide functionally involved in the DNA mismatch repair system of a plant, for example a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human such as a polypeptide encoded by the genes AtMSH3 or AtMSH6 of Arabidopsis thaliana, as defined hereinbelow.

According to a third embodiment of the invention, there is provided an isolated and purified DNA molecule comprising a polynucleotide sequence selected from the group consisting of (i) a sequence encoding a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence; and (ii) a sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant.

According to a fourth embodiment of the invention there is provided a chimeric gene comprising a DNA sequence selected from the group consisting of (i) a sequence encoding a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence, and (ii) a sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant: together with at least one regulation element capable of functioning in a plant cell. Examples of such regulation elements include constitutive, inducible, tissue type specific and cell type specific promoters such as 35S, NOS, PR1a, AoPR1 and DMC1. Typically, a chimeric gene of the fourth embodiment will also include at least one terminator sequence, more typically exactly one terminator sequence.

In the third and fourth embodiments, said interference, by said polynucleotide sequence, with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair peptide of a yeast or a human typically occurs by hybridisation or by co-suppression.

According to a fifth embodiment of the invention there is provided a plasmid or vector comprising a chimeric gene of the fourth embodiment. A vector of the fifth 30 embodiment may be, for example, a viral vector or a bacterial vector.

According to a sixth embodiment of the invention, there is provided a plant cell stably transformed, transfected or electroporated with a plasmid or vector of the fifth embodiment.

According to seventh embodiment of the invention, there is provided a plant 35 comprising a cell of the sixth embodiment.

According to an eighth embodiment of the invention, there is provided a process for at least partially inactivating a DNA mismatch repair system of a plant cell, comprising

transforming or transferring said plant cell with a DNA sequence of the third embodiment or a chimeric gene of the fourth embodiment or a plasmid or vector of the fifth embodiment, and causing said DNA sequence to express said polynucleotide or said polypeptide.

According to a ninth embodiment of the invention, there is provided a process for increasing genetic variation in a plant comprising obtaining 'a hybrid plant from a first plant and a second plant, or cells thereof, said first and second plants being genetically different; altering the mismatch repair system in said hybrid plant; permitting said hybrid plant to self-fertilise and produce offspring plants; and screening said offspring plants for plants in which homeologous recombination has occurred. For example, homeologous recombination may be evidenced by new genetic linkage of a desired characteristic trait or of a gene which contributes to a desired characteristic trait.

According to a tenth embodiment of the invention there is provided a process for obtaining a plant having a desired characteristic, comprising altering the mismatch repair system in a plant, cell or plurality of cells of a plant which does not have said desired characteristic, permitting mutations to persist in said cells to produce mutated plant cells, deriving plants from said mutated plant cells, and screening said plants for a plant having said desired characteristic.

In a preferred form of the ninth and tenth embodiments of the invention, the step of altering the mismatch repair system comprises introducing into said hybrid plant, plant, cell or cells a chimeric gene of the fourth embodiment and permitting the chimeric gene to express a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence in a mismatch repair gene of the hybrid plant, plant, cell or cells, or a polypeptide capable of disrupting the DNA mismatch repair system of the hybrid plant or cells.

In other embodiments, the invention provides (a) an oligonucleotide capable of hybridising at 45°C under standard PCR conditions to a DNA molecule of the first embodiment; (b) an oligonucleotide capable of hybridising at 45°C under standard PCR conditions to the DNA of SEQ ID NO: 18 and (c) an oligonucleotide capable of hybridising at 45°C under standard PCR conditions to the DNA of SEQ ID NO:30; with the proviso that the oligonucleotide of (a), (b) and (c) is other than SEQ ID NO:1 or SEQ ID NO:2.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides a diagrammatic representation of the primer sequences used to 35 isolate AtMSH3.

Figure 2 is a plasmid map of clone 52, showing restriction enzyme cleavage sites in the 5' half of the full-length cDNA for AtMSH3.

Figure 3 is a plasmid map of clone 13, showing restriction enzyme cleavage sites in the 3' half of the full-length cDNA for AtMSH3.

Figure 4 is a sequence listing of the coding sequence of AtMSH3, together with a deduced sequence of the encoded polypeptide.

Figure 5 is a protein alignment of yeast (Saccharomyces cerevisiae) and Arabidopsis thaliana MSH3 protein.

Figure 6 provides a diagrammatic representation of the primer sequences used to isolate AtMSH6.

Figure 7 is a plasmid map of clone 43, showing restriction enzyme cleavage sites in the 5' half of the full-length cDNA for AtMSH6.

Figure 8 is a plasmid map of clone 62, showing restriction enzyme cleavage sites in the 3' half of the full-length cDNA for AtMSH6.

Figure 9 is a sequence listing of the coding sequence of AtMSH6, together with a deduced sequence of the encoded polypeptide.

Figure 10 is a protein alignment of yeast (Saccharomyces cerevisiae) and Arabidopsis thaliana MSH6 protein.

Figure 11 is a genomic sequence listing of AtMSH6.

Figure 12 is a plasmid map of plasmid pPF13.

Figure 13 is a plasmid map of plasmid pPF14.

Figure 14 is a plasmid map of plasmid pCW186.

Figure 15 is a plasmid map of plasmid pCW187.

Figure 16 is a plasmid map of plasmid pPF66.

Figure 17 is a plasmid map of plasmid pPF57.

Figure 18 is a diagrammatic representation of an antisense gene construction for use in homeologous meiotic recombination.

Figure 19 is a plasmid map of plasmid p3243.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is based on the inventors' discovery that there exist in higher plants genes which are homologous to MMR genes in E. coli, and to MMR genes in yeasts and humans.

Thus, the inventors have identified genes, herein designated AtMSH3 and AtMSH6, of the plant Arabidopsis thaliana which encode the proteins AtMSH3 and AtMSH6. These plant proteins are homologous to yMSH3 and yMSH6, respectively. The present inventors have isolated cDNAs encoding the proteins AtMSH3 and AtMSH6 and have isolated the complete gene encoding AtMSH6. Given the teaching herein, other genes (for example AtMSH2, and genes of other plants) may be obtained which are involved in DNA mismatch repair in plants, including other genes which encode polypeptides homologous to MMR proteins of yeasts or humans, such as genes which encode

polypeptides homologous to yeast MSH2. MLH1 or PMS2, or to human MLH1. PMS1 or PMS2. For example, given the teaching herein, genes of members of the *Brassicaceae* family or of other unrelated families, for example the *Poaceae*, the *Solanaceae*, the *Asteraceae*, the *Malvaceae*, the *Fabaceae*, the *Linaceae*, the *Canabinaceae*, the *Dauaceae* and the *Cucurbitaceae* family, and which encode polypeptides homologous to MMR proteins of yeasts or humans may be obtained.

Examples of plants whose genes encoding polypeptides homologous to MMR proteins of yeasts or humans may be obtained given the teaching herein include maize, wheat, oats, barley, rice, tomato, potato, tobacco, capsicum, sunflower, lettuce, artichoke, safflower, cotton, okra, beans of many kinds including soybean, peas, melon, squash, cucumber, oilseed rape, broccoli, cauliflower, cabbage, flax, hemp, hops and carrot.

Within the meaning of the present invention, a first polypeptide is defined as homologous to a second polypeptide if the amino acid sequence of the first polypeptide exhibits a similarity of at least 50% on the polypeptide level to the amino acid sequence of the second polypeptide.

A procedure which may be followed to obtain genes AtMSH3 and AtMSH6 is described in Example 1. Essentially the same technique may be applied to obtain other mismatch repair genes of Arabidopsis thaliana, and essentially the same technique as 20 exemplified herein may be applied to cDNA obtained by reverse transcription of RNA from other plants. Alternatively, given the sequence information disclosed herein, other degenerate oligonucleotide primers, especially oligonucleotides of the invention which are capable of hybridising at 45°C under standard PCR conditions (such as the conditions described in Example 1 using primers UPMU and DOMU) to AtMSH3 and/or AtMSH6 25 may be designed and obtained for use in isolating sequences of plant mismatch repair genes which are homologous to AtMSH3 or AtMSH6, from other plants. oligonucleotides of the invention which are capable of hybridising at 45°C under standard PCR conditions to plant mismatch repair genes of plants other than Arabidopsis thaliana also fall within the scope of the present invention and may be utilised to obtain mismatch Typically, such oligonucleotides are capable of 30 repair genes of still other plants. hybridising at 45°C under standard PCR conditions to a DNA molecule which encodes a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or a human. The temperature at which oligonucleotides of the invention hybridise to AtMSH3 and/or AtMSH6, or to plant mismatch repair genes of plants other than Arabidopsis thaliana, or 35 to DNA molecules which encode polypeptides which are homologous to a mismatch repair polypeptide of a yeast or a human may be higher than 45°C, for example at least 50°C, or at least 55°C, or at least 60°C or as high as 65°C.

The successful gene isolation disclosed herein demonstrates for the first time the existence of MMR in higher plants and indicates the presence of other plant MMR genes. For example, genes encoding the plant homologs of MSH1, MSH2, MSH4, MSH5, PMS1, PMS2 and MLH1 may be identified given the teaching herein. Such genes, as well as those specifically described herein, separately or in combination, are useful in manipulating the plant MMR for plant breeding purposes. Thus, for example, the plant MMR may be altered by including in a plant cell a polynucleotide sequence as defined herein above with reference to the third embodiment of the invention, and causing the polynucleotide sequence to express either a polynucleotide which disables a plant MMR gene, or a polypeptide which disrupts the plant's MMR system.

The DNA molecule of the third embodiment of the invention includes a polynucleotide sequence (herein referred to as a MMR altering gene) which may for example encode sense, antisense or ribozyme molecules characterised by sufficient base sequence similarity or complementarity to the gene to be altered to permit the antisense or 15 ribozyme molecule to hybridise with the plant MMR gene in vivo or to permit the sense molecule to participate in co-suppression. Alternatively, the MMR altering gene may encode a protein or proteins which interfere with the activity of a plant MMR protein and thus disrupt the plant's MMR system. For example, such encoded proteins may be antibodies or other proteins capable of interfering with MMR protein function, such as by 20 complexing with a protein functionally involved in plant MMR thereby disrupting the MMR of the plant. An example of such a protein is the MSH3 protein of Arabidopsis thaliana described herein or a protein of another plant which is homologous to the MSH3 protein of A. thaliana. For instance, overexpression of MSH3 in a plant cell causes MSH2 present in the cell to be substantially completely complexed, disrupting the 25 mismatch repair mechanism or mechanisms in the cell which are functionally dependent on the presence of a complex of MSH2 with MSH6. Similarly, mismatch repair mechanisms which depend on the presence of a complex of MSH2 and MSH3 may be disrupted by the overexpression of MSH6.

A chimeric gene of the fourth embodiment, incorporating a MMR altering gene, may be prepared by methods which are known in the art. Similarly, the MMR altering gene may be introduced into a plant cell, regenerating tissue or whole plant by techniques known in the art as being suitable for plant transformation, or by crossing. Known transformation techniques include Agrobacterium tumefaciens or A. rhizogenes mediated gene transfer, ballistic and chemical methods, and electroporation of protoplasts.

The MMR altering gene or genes are typically expressed from suitable promoters. Suitable promoters may direct constitutive expression, such as the 35S or the NOS promoter. Usually, however, the promoter will direct either inducible or tissue specific (e.g. callus; embryonic tissue; etc.), cell type specific (e.g. protoplasts: meiocytes; etc.) or developmental (e.g. embryo) expression of the altering gene or genes, in order for the

WO 99/19492 PCT/EP98/06977

MMR system to function in tissue types or cell types, or at developmental stages of the plant, in which it is not desirable for the MMR system to be altered. Using such promoters, therefore, the activity of a MMR altering gene may be limited to a specific stage during plant development or it may be altered by controlling conditions external to the plant, and the deleterious effects of a permanently disabled or altered DNA mismatch repair system in a plant may be avoided. Examples of suitable promoters which are not constitutive are known in the art and include inducible promoters such as *PR*1a (reviewed by Gatz, 1997, Annual Rev. Plant Phys. Plant Mol. Biol. 48: 89), tissue specific promoters such as *AoPR1* (Sabahattin et al., 1993, Biotechnology 11: 218), and cell-type specific promoters such as *DMC*1.

A chimeric gene in accordance with the invention may further be physically linked to one or more selection markers such as genes which confer phenotypic traits such as herbicide resistance, antibiotic resistance or disease resistance, or which confer some other recognisable trait such as male sterility, male fertility, grain size, colour, growth rate, flowering time, ripening time, etc.

The process of the tenth embodiment of the invention provides, for example, a process for generating intraspecies genetic variation by altering the mismatch repair system in a plant cell, in regenerating plant tissue or in a whole plant. The plant cell, regenerating tissue or whole plant includes and expresses one or more MMR altering 20 genes which are capable of altering mismatch repair in the plant cell, regenerating tissue or whole plant. Alteration of MMR may be achieved, for example, by inactivating the genes encoding plant MSH3 and/or plant MSH6. It is preferred to inactivate the plant MSH3 and MSH6 encoding genes at the same time and in the same plant cell, regenerating tissue or whole plant. Typically in this preferred form of the invention 25 inactivation of either plant MSH3 or MSH6 alone is insufficient to substantially alter the plant's mismatch repair system and only when both MSH3 and MSH6 are inactivated simultaneously is the plant's mismatch repair system sufficiently altered to prevent the MMR system from recognising base pair mismatches, base insertions or deletions as a result of DNA replication errors, DNA damage, or oligonucleotide induced site-specific 30 mutagenesis. However, in some applications of the invention, inactivation of only one gene may also be used to cause genomic instability or increase the efficiency of sitespecific mutagenesis.

If desired, the MMR altering gene or genes may later be rendered non-functional or ineffective, or may be removed from the genome of the plant cell, regenerating tissue or whole plant in order to restore mismatch repair in the plant cell, regenerating tissue or whole plant. The MMR altering gene or genes may be inactivated by means of known gene inactivation tools, such as ribozymes, or may be removed from the genome using gene elimination systems known in the art, such as *CRE/LOX*. It is preferred to render two genes, whose gene products combine to incapacitate MMR, ineffective by separating

the altering genes through segregation. Therefore, in a preferred embodiment of the invention a first plant cell or plant is generated in which only plant MSH3 is incapacitated, and a second plant cell or plant is generated in which only plant MSH6 is incapacitated. The combination of both genomes, for example by crossing, then produces significant 5 MMR deficiency in those cells or plants which have inherited both altering genes. If the altering genes are expressed from unlinked loci, gene segregation restores MMR activity in the progeny of the cells or plants.

In a process of the ninth embodiment of this invention, homeologous recombination is enhanced between different genomes, chromosomes or genes in plant cells or plants by altering MMR in said plant cells or plants. Such genomes, chromosomes or genes are characterised in that they originate from different plant families, genera, species, subspecies, plant varieties or lines. Hybrid plant cells or hybrid plants may be produced by crossing, by cell fusion or by other techniques known in the art. These plant cells or plants are further characterised by expressing one or more genes that are capable of altering mismatch repair in the plant cell or plants.

In the process of the ninth embodiment, the homeologous recombination is typically for the purpose of introducing a desired characteristic in the hybrid plant. In this typical application of the process of the ninth embodiment, and in the process of the tenth embodiment the desired characteristic may be any characteristic which is of value to the plant breeder. Examples of such characteristics are well known in the art and include altered composition or quality of leaf or seed derived storage products (e.g. oil, starch, protein), altered composition or quality of cell walls (e.g. decrease in lignin content), altered growth rate, prolonged flowering, increased plant yield or grain yield, altered plant morphology, resistence to pathogens, tolerance to or improved performance under environmental stresses of various kinds, etc.

In a preferred form of the tenth embodiment, an MMR altering gene is cointroduced along with the homeologous genome, chromosome or gene of another plant
cell or plant into an MMR proficient plant cell or MMR proficient plant to produce a
hybrid plant cell or hybrid plant in which homeologous recombination can occur.

Suitably, the MMR proficient plant cell or MMR proficient plant may also include an
MMR altering gene. For example a gene capable of inactivating plant MSH3 may be cointroduced along with the homeologous genome, chromosome or gene of another plant
cell or plant into an MMR proficient plant cell or MMR proficient plant in which MSH6
is inactivated. A resultant hybrid plant in which homeologous recombination occurs will
include both the MSH3 and MSH6 altering genes and its MMR system will therefore be
inactivated.

In this form of the invention, if hybrid plants are to be produced by crossing, the MMR altering gene preferably originates from the male parent, thus ensuring that the

MMR altering gene is always introduced and is not present in the recipient cell. That is, the MMR of the recipient cell, prior to introduction of the MMR altering gene, is typically proficient. Alternatively, if an MMR altering gene is present in a recipient cell it may be ineffective or inefficient on its own, or it may be linked to an inducible or tissue specific or cell type specific promoter which only renders the MMR altering gene active under limited conditions.

Thus, in a preferred form of the process of the ninth embodiment, the MMR system of the hybrid plant is initially unaltered. In this form of the process, the step of altering the mismatch repair system may comprise introducing into the hybrid plant, or cells thereof, a MMR altering gene, such as by Agrobacterium tumefaciens or A. rhizogenes mediated gene transfer, ballistic and chemical methods, and electroporation of protoplasts.

The MMR altering gene or genes are typically expressed from suitable promoters, as described above. Preferably, the promoter is transcriptionally active in mitotically and meiotically active tissue and/or cells to ensure MMR alteration after chromosome pairing at mitosis and meiosis, respectively. The preferred timing for MMR alteration is at meiosis, because recombinant genomes, chromosomes or genes are directly transmitted to the progeny. A suitable meiocyte specific promoter is for example the *DMC*1 promoter from *Arabidopsis thaliana* ssp. *Ler.* (Klimyuk and Jones, 1997, Plant J. 11, 1-14). However, mitotic homeologous recombination is also a desirable outcome as somatic recombination events can be transmitted to offspring due to the totipotency of plant cells and the lack of predetermined germ cells in plants.

If desired, the MMR altering gene or genes may later be rendered non-functional or ineffective, or may be removed from the hybrid plant or hybrid plant cells, in order to restore mismatch repair in the hybrid plant or hybrid plant cells. The MMR altering gene or genes may be inactivated by means of known gene inactivation tools as described herein above.

EXAMPLES

Example 1. Cloning of the AtMSH3 and AtMSH6 coding sequences

Isolation of partial AtMSH3 and AtMSH6 consensus sequences

30

Degenerate oligonucleotides UPMU (SEQ ID NO:1) and DOMU (SEQ ID NO:2)

UPMU CTGGATCCACIGGICCIAA(C/T)ATG

DOMU CTGGATCC(A/G)TA(A/G)TGIGTI(A/G)C(A/G)AA

were used to isolate AtMSH3 and AtMSH6 sequences by PCR amplification.

Primers UPMU and DOMU correspond to conserved amino acid sequences of the proteins MutS (E. coli and S. typhimurium), HexA (S. pneumoniae). Rep! (mouse) and Ducl (human). The conserved regions to which they are targeted are TGPNM for UPMU (amino acid positions 852-856 for AtMSH6 and 816-820 for AtMSH3) FATHY or FVTHY

for DOMU (amino acid positions 964-968 for AtMSH6 and 928-932 for AtMSH3, respectively.) These primers have been used to isolate MSH2 and MSH1 from yeast (Reenan and Kolodner, Genetics 132: 963-973 (1992)) and MSH2 from *Xenopus* and mouse (Varlet et al., Nuc. Acids Res. 22:5723-5728 (1994)).

Template single strand cDNA was produced by reverse transcription of 2 µg total RNA from a cell suspension culture of Arabidopsis thaliana ecotype Columbia (Axelos et al. 1989, Mol. Gen. Genetics 219: 106-112). The PCR reaction was performed under the following conditions in a final volume of 100µl: 0.2mM dNTP, 1µM each primer, IXPCR buffer, 1u Taq DNA polymerase (Appligene) in the presence of template cDNA. PCR 10 parameters were 5 minutes at 94°C, followed by 30 cycles of 40 seconds at 95°C, 90 seconds at 45°C, 1 minute at 72°C. The amplification products were cloned into pGEM-T vector (Promega) and sequenced. Two different clones were isolated, S5 (350bp) was homologous to MSH3, S8 (327bp) was homologous to MSH6. Complete cDNA sequences were then isolated according to the Marathon cDNA amplification kit procedure (Clontech). 15 In summary, this procedure involves producing double stranded cDNA by reverse transcription of 2µg polyA+ RNA from the cell suspension culture of Arabidopsis. Adaptors are ligated on each side of the cDNA. The ligated cDNA is used as a template for 5' and 3' RACE PCR reactions in the presence of primers that are specific for the adaptor on one side (AP1 and AP2), and specific for the targeted gene on the other side. A 5' and a 3' 20 fragment that overlap are thus produced for each gene. The complete gene coding sequence can be reconstituted taking advantage of a unique restriction site, if available, in the overlapping region. Specific details of this procedure as it was used to isolate AtMSH3 and AtMSH6 coding regions, are as follows.

Isolation of AtMSH3 complete coding sequence

25

From the sequence of clone S5, primer 636 (SEQ ID NO:3) was designed:

636 TGCTAGTGCCTCTTGCAAGCTCAT.

Primer AP1 (SEQ ID NO:4) is complementary to a portion of an adaptor sequence which had been ligated to the 5' and 3' ends of Arabidopsis cDNA:

API CCATCCTAATACGACTCACTATAGGGC.

PCR performed on the ligated cDNA with primers 636 and AP1 for the 5' RACE PCR was followed by a second round of amplification with the nested primers AP2 (SEQ ID NO:5) and S525 (SEQ ID NO:6)

AP2 ACTCACTATAGGGCTCGAGCGGC

S525 AGGTTCTGATTATGTGTGACGCTTTACTTA

35 (the latter was also designed to correspond to a part of the sequence of clone S5) and produced a 2720bp DNA fragment. Figure 1 provides a diagrammatic representation of the primer sequences used to isolate AtMSH3. Another primer (S51, SEQ ID NO:7)

S51 GGATCGGGTACTGGGTTTTGAGTGTGAGG

was designed closer to the 5° border and permitted the determination of 99bp upstream to the ATG initiation codon. For the 3° RACE PCR, a first PCR reaction was performed with primers AP4 and 635 (SEQ ID NO:8).

- 635 GCACGTGCTTGATGGTGTTTTCAC
- 5 followed by a second round of amplification, using the nested primers AP2 and S523 (SEQ ID NO:9)
 - S523 TCAGACAGTATCCAGCATGGCAGAAGTA

which produced a DNA fragment of 890bp. Both DNA fragments were subcloned into pGEM-T and sequenced. Since PCR amplification using the Expand Long Template PCR system (Boehringer-Mannheim) produced errors in the sequence, new oligonucleotides were designed to isolate those sequences again by PCR, but with the high fidelity DNA polymerase Pfu. PCR with primers 1S5 (SEQ ID NO:10) and S53 (SEQ ID NO:11)

- 1S5 ATCCCGGGATGGGCAAGCAAAAGCAGCAGACGA
- S53 GACAAAGAGCGAAATGAGGCCCCTTGG
- amplified the 1244bp fragment clone 52 (SEQ ID NO:12, cloned into pUC18/Sma1). PCR with primers S52 (SEQ ID NO:13) and 2S5 (SEQ ID NO:14)
 - 2S5 ATCCCGGGTCAAAATGAACAAGTTGGTTTTAGTC
 - S52 GCCACATCTGACTGTTCAAGCCCTCGC

amplified the 2104bp clone 13 (SEQ ID NO:15, cloned into pUC18/Smal). The complete coding sequence of the AtMSH3 gene was reconstructed in pUC18 by ligating the 5' half of AtMSH3 (clone 52) to the 3' half of AtMSH3 (clone 13) after digesting with BamH1 which has a unique cleavage site in the overlapping region of both clones. This manipulation yielded plasmid pPF26. The Smal fragment from pPF26 contains the complete AtMSH3 coding sequence. The remaining primers referred to in Figure 1 are as follows:

- S51 GGATCGGGTACTGGGTTTTGAGTGTGAGG (SEQ ID NO:16)
- S525 AGGTTCTGATTATGTGTGACGCTTTACTTA (SEQ ID NO:17)

Figures 2 and 3 provide plasmid maps of clones 52 and 13 respectively, showing restriction enzyme cleavage sites. The complete AtMSH3 coding sequence (SEQ ID NO:18) 30 is 3246bp long and is shown in Figure 4 together with the deduced sequence (SEQ ID NO:19) of the encoded polypeptide. AtMSH3 is clearly homologous to the yeast and mouse MSH3 genes. A sequence alignment of polypeptides encoded by AtMSH3 and that encoded by Saccharomyces cerevisiae MSH3 is set out in Figure 5.

Isolation of the AtMSH6 complete coding sequence and genomic sequences

The same procedure allowed isolation of the AtMSH6 cDNA. Figure 6 provides a diagrammatic representation of the primer sequences used to isolate AtMSH6. For the 5' RACE PCR, primers 638 (SEQ ID NO:20) and AP1 (SEQ ID NO:4)

638 TCTCTACCAGGTGACGAAAAACCG
allowed the amplification of a 2889 DNA fragment. Primer S81 (SEQ ID NO:21)

S81 CGTCGCCTTTAGCATCCCCTTCCTTCAC

helped define the 142bp upstream to the ATG initiation codon. On the 3' side, RACE PCR was initially performed with primers S823 (SEQ ID NO:22) and API (SEQ ID NO:4),

S823 GCTTGGCGCATCTAATAGAATCATGACAGG

5 and then with the nested primers 637 (SEQ ID NO:23) and AP2 (SEQ ID NO:5).

637 GACAGCGTCAGTTCTTCAGAATGC

to produce a 774bp DNA fragment. As for *AtMSH3*, those fragments were cloned and sequenced. Re-isolation of the DNA sequence using the high fidelity *Pfu* polymerase and newly designed primers 1S8 (SEQ ID NO:24) and S83 (SEQ ID NO:25) (for the 5' side) led to a 2182 bp DNA fragment identified as clone 43 (SEQ ID NO:26, cloned in pUC18/Sma1), and a 1379bp clone identified as clone 62 (SEQ ID NO:27, also cloned in pUC18/Sma1).

- 1S8 ATCCCGGGATGCAGCGCCAGAGATCGATTTTGT
 2S8 ATCCCGGGTTATTTGGGAACACAGTAAGAGGATT (SEQ ID
 NO:28)
 S82 GCGTTCGATCATCAGCCTCTGTTGC (SEQ ID NO:29)
 - S83 CGCTATCTATGGCTGCTTCGAATTGAG

Figures 7 and 8 provide plasmid maps of clones 43 and 62 respectively, showing restriction enzyme cleavage sites. Clones 43 and 62 were digested by the Xmn1 restriction enzyme for which a unique site is present in their overlapping region and then ligated. The complete AtMSH6 coding sequence (SEQ ID NO:30) is 3330bp long and is shown in Figure 9 together with the deduced sequence (SEQ ID NO:31) of the encoded polypeptide. AtMSH6 is clearly homologous to the yeast and mouse MSH6genes. A sequence alignment of polypeptides encoded by AtMSH6 and that encoded by Saccharomyces cerevisiae MSH6 is set out in Figure 10.

An AtMSH6 genomic sequence was also isolated from a genomic DNA library constituted after partial Sau3Al digestion of DNA from the Arabidopsis cell suspension. 8062bp were sequenced that covered the AtMSH6 gene and show colinearity with the cDNA. 16 introns are found scattered along the gene. The complete genomic sequence 30 (SEQ ID NO:98) is shown in Figure 11.

Example 2. A measure of somatic variation in MMR deficient plants Constructs

Constructs with antisense AtMSH3 or antisense AtMSH6 or both AtMSH3/AtMSH6 under the control of a single 35S promoter have been inserted into the binary vector pPZP121 (Hajdukiewicz et al., Plant Mol. Biol. 23, 793-799) between the right and left borders of the T-DNA. The pPZP121 plasmid confers chloramphenical resistance to Escherichia coli or Agrobacterium tumefaciens bacteria. The aacC1 gene is carried by the T-DNA and allows selection of transformed plant cells on gentamycin (Hajdukiewicz et al., Plant Mol. Biol. 25, 989-994). For the purpose of expressing antisense constructs, a 35S

promoter/terminator cassette with a polylinker was introduced into pPZP121. The 3' ends of the considered genes have been chosen since this region seems more efficient for antisense inhibition. For AtMSH3 this corresponds to clone 13 (2104bp), for AtMSH6 this corresponds to clone 62 (1379bp). Clone 13 comprises 2104bp of the 3' region that were cut off the pUC18 vector by Sal1/Sst1 restriction, blunted with T4 DNA polymerase and ligated into the T4 DNA polymerase blunted BamHI site of pPZP121/35S, creating clone pPF13. The same procedure was followed for the 3' region of AtMSH6 clone 62 (1379bp) thus creating plasmid pPF14. For the double constructs, the 3' region (from clone 62) of AtMSH6 was introduced ahead of the AtMSH3 region into pPF13 creating pCW186 and reciprocally, the 3' region of AtMSH3 (from clone 13) was introduced ahead of AtMSH6 into pPF14, creating pCW187.

These constructs were introduced into the Arabidopsis cells (as described below) of wildtype Columbia and of the Columbia tester line.

An alternative strategy to antisense inhibition of AtMSH6 comes from experiments of Marra et al. (1998, Proc. Natl. Acad. Sci USA 95, 8568-8573) who show that overexpression of functional MSH3 results in depletion of MSH6 protein in human cells. This depletion may generate a mismatch repair mutant phenotype.

For the purpose of overexpressing functional AtMSH3 protein in plant cells, the complete MSH3 coding region was excised from pPF26 (example 1) by digestion with 20 Smal, and was inserted into the Smal site of pCW164. The resulting construct was named pPF66. It contains a complete AtMSH3 gene under the control of the 35S promoter inside the left (LB) and right (RB) border of the T-DNA. This T-DNA also contains the hpt2 gene for gentamycin selection. Plasmid pPF66 was introduced into Arabidopsis cells as described below. One cell clone was selected which clearly overexpressed the AtMSH3 gene as shown by Northern analysis. Figures 12-16 provide plasmid maps of plasmids pPF13, pPF14, pCW186, pCW187 and pPF66, respectively.

Construction of tester construct

For the purpose of Forward Mutagenesis Assays, a tester construct was built containing the coding regions for nptII, codA, uidA. All three genes are driven by the 35S promoter and are terminated by the 35S terminator. This construct was obtained by introducing an EcoR1 fragment encoding the codA cassette (2.5kb) and a HindIII fragment encoding the uidA (GUS) cassette (2.4kb) into the pPZP111 vector (Hajdukiewicz et al.,1994, Plant Mol Biol 23: 793-799) which already contained the nptII expression cassette. This new plasmid was named pPF57. NptII is used to select for transformed plant cells. 35 GUS is used to analyse the degree of gene silencing in the construct (i.e. to identify cell lines in which the transgenes are expressed), and codA is used as a marker for forward mutagenesis (described below).

The plasmid map of pPF57 is provided in Figure 17.

Plant cell transformation

then transformed by co-cultivation. A suspension culture of Arabidopsis thaliana cells that 5 has been established by Axelos et al. (1992, Plant Physiol. Biochem. 30, 1-6) may be used. One day old freshly subcultured cells are diluted five times into AT medium (Gamborg B5 medium. 30g/l sucrose, 200µg/l NAA). 10µl of saturated Agrobacterium containing the transforming T-DNA constructs are added to 10ml diluted cells in a 100ml erlenmeyer. The co-cultivation is agitated slowly (80rpm) for 2 days. The cells are then washed 3 times into AT medium and finally resuspended in the same initial volume (10ml). The culture is agitated for 3 days to allow expression before plating on selection plates (AT/BactoAgar 8g/l+gentamycin 50µg/ml). Transformed individual calli are isolated 3 weeks later.

Tester Strain

The tester construct on plasmid pPF57 was introduced into Arabidopsis cells of wildtype Columbia using the transformation protocol described above. Among 10 candidate transformants, one cell clone was shown (by Southern analysis) to have a unique T-DNA insertion. All three genes were shown to be functional in this cell line as indicated by resistance to kanamycin, blue staining in the presence of X-Glu (GUS), and sensitivity to 5-fluoro-cytosine (codA).

MMR altering genes (described above) were then introduced individually into the tester line and transformed cells are used for analysis of both Microsatellite Instability and Forward Mutagenesis.

Microsatellite analysis

Microsatellites have been described in Arabidopsis (Bell and Ecker, 1994, Genomics 19, 137-144). The present Example is based on a study of instability of microsatellites that are polymorphic amongst different ecotypes. DNA is extracted from the transformed calli. Specific primers have been defined that are used to amplify the microsatellite sequence. One of the two primers is previously P³² labelled by T4 kinase. In case of a polymorphic variation, new PCR products appear that do not follow the expected pattern of migration on a polyacrylamide gel. This is a commonly observed feature for MMR deficient cells in yeast or mammalian cells.

In particular, the present Example describes a study on microsatellites ca72 (CA₁₈), nga172 (GA₂₉), and ATHGENEA(A₃₉), chosen because they belong to the types predominantly affected in human mismatch repair deficient tumors. The size of these microsatellites is not conserved from one *Arabidopsis* ecotype to the other.

Arabidopsis cells which are transformed with an MMR altering gene (above) and control cells not expressing the MMR altering gene are allowed to form calli. DNA is

rapidly extracted from the calli and is analysed for microsatellite instability as described in detail by Bell and Ecker 1994. Genomics 19, 137-144. In summary, the relevant microsatellite is amplified by PCR using P32 labelled primers. The PCR products are separated on a DNA sequencing gel for size determination. Size differences between microsatellites from transformed and control cells not expressing the MMR altering gene in question indicate microsatellite instability as a result of MMR alteration.

The sequences of primers used for PCR amplification of microsatellites ca72 and nga172 are included in Table 1. PCR amplification of microsatellite ATHGENEA made use of a forward primer containing the sequence

ACCATGCATAGCTTAAACTTCTTG (SEQ ID NO:32)

and of a reverse primer containing the sequence

ACATAACCACAAATAGGGGTGC (SEQ ID NO:33).

The amplification for microsatellite ca72 revealed in Columbia control cells (with respect to the MMR altering gene) a 248 bp long PCR fragment instead of the published length of 124 bp. DNA sequencing verified this fragment as a CA₁₈ microsatellite.

Forward mutagenesis assay

10

Tester cells transformed with antisense AtMSH3 or antisense AtMSH6 or both AtMSH3/AtMSH6 are analysed for the stability of the codA gene. The functional codA gene confers to sensitivity to 5-fluoro-cytosine (5FC), whereas a gene inactivating mutation in codA will confer resistance to 5FC. The frequency of resistant cells is therefore a good indicator of somatic variation as a direct result of MMR alteration. Variants resistant to 5FC are first analysed for GUS activity. If GUS is inactive, 5FC resistance is assumed to be due to gene silencing (all three genes are under the 35S promoter). If GUS is active, 5FC resistance is assumed to be due to forward mutations that have inactivated codA. PCR is then performed on the putative codA mutant genes which is then sequenced to confirm the presence of forward mutations in codA.

Besides *codA*, other marker genes may also be used for the Forward Mutagenesis Assay such as the *ALS* gene (conferring sensitivity to valine or to sulfonylurea; Hervieu and Vaucheret, 1996, Mol. Gen. Genet. 251 220-224; Mazur et al. 1987, Plant Physiol. 85 1110-30 1117).

Example 3. Homeologous meiotic recombination in Arabidopsis thaliana

A. Construction of a melocyte specific gene expression cassette comprising the DMC1 promoter and the NOS terminator

(i) The DMC1 promoter may be used as published by Klimyuk and Jones, 1997, 35 Plant J. 11.1-14). To obtain a more convenient alternative for gene cloning, a 3.3 Kb

long subfragment of the DMC1 promoter was obtained by PCR from genomic DNA of Arabidopsis thaliana (ssp. Landsberg erecta "Ler").

The PCR was done in three rounds:

10

Round One: A 3.7 Kb long product was obtained using the forward primer 5 DMCIN-A comprising the sequence

GAAGCGATATTGTTCGTG (SEQ ID NO:34)

and the reverse primer DMCIN-B comprising the sequence

AGATTGCGAGAACATTCC (SEQ ID NO:35).

The weak amplification product was then used as template for round two and three.

Round Two: A 3.1 Kb long product comprising the promoter and the 5' untranslated leader was obtained using forward primer DMCIN-1, which contained the sequence

acgcgtcgacTCAGCTATGAGATTACTCGTG (SEQ ID NO:36)

and introduced a Sall cloning site at the 5' end of the promoter fragment, and reverse primer DMCIN-2 which contained the sequence

gctctagaTTTCTCGCTCTAAGACTCTCT (SEQ ID NO:37)

and introduced a XbaI site at the 3' end of the PCR fragment.

Round Three: A 0.2 Kb long product comprising the first exon/intron of the *DMC*1 promoter was obtained using forward primer DMCIN-3, which contained the sequence

gctctagaGCTTCTCTTAAGTAAGTGATTGAT (SEQ ID NO:38)

and introduced a XbaI site at the 5' end of the PCR fragment, and reverse primer DMCIN-4, containing the sequence

teccegggetegagagatetecatggTTTCTTCAGCTCTATGAATCC (SEQ ID NO:39) and introduced at the 3' end of the PCR product restriction sites for Ncol, Bg/II. XhoI and 25 SmaI.

The products obtained in round Two and Three were digested with XbaI and subsequently ligated to reconstitute a 3.3 Kb long DMC1 promoter from which the first two in-frame ATG start codons were replaced with a unique restriction site for XbaI. This promoter can be cloned between the restriction sites for SalI and SmaI of p3264, which contains the SacI-EcoRI NOS terminator in pBIN19, to yield the entire expression cassette in pBIN19. This cassette contains the following cloning sites: NcoI, BglII, XhoI. SmaI and (already present on p3264) KpnI and SacI.

(ii) Another strategy yielded the following convenient *DMC*1 promoter. A 1.8 kb DNA fragment comprising the 3' terminal part of the melocyte specific *DMC*1 promoter was isolated by PCR from purified genomic DNA of *Arabidopsis thaliana* (ssp. Landsberg erecta "Ler"). The forward PCR primer (DMC1a) contained the sequence

acgcgtcgacGAATTCGCAAGTGGGG (SEQ ID NO:40)

and introduced a Sall cloning site at the 5' end of the promoter fragment. The reverse PCR primer (DMC1b) contained the sequence

tecatggagatetecegggtaeCGATTTGCTTCGAGGG (SEQ ID NO:41) introducing a polylinker region at the 3' end of the promoter fragment. The PCR reaction was carried out with VENT DNA Polymerase (NEB) over 25 cycles using the following cycling protocol: 1 minute at 94°C, 1 minute at 54°C, 2 minutes at 72°C.

The PCR reaction yielded a blunt ended DNA fragment which was digested with restriction enzyme SaII and was cloned into the cleavage sites of restriction enzymes SaII and SmaI in plasmid p2030, a pUC118 derivative containing the SacI-EcoRI NOS terminator fragment of pBIN121. The cloning yielded plasmid p2031, containing the DMC1 promoter-polylinker-NOS terminator expression cassette depicted in Figure 18.

10 B. Construction of an MSH3 antisense gene under the control of the DMC1 promoter

A 2.1 kb DNA fragment encoding the carboxyterminal part of AtMSH3 was removed from the polylinker of clone 13 described in Example 1 after (i) digestion with KpnI. (ii) blunting of the DNA ends generated by KpnI and (iii) digestion with BamHI. The isolated fragment was then cloned in antisense orientation downstream of the DMC1 promoter in plasmid p2031, which had been digested with restriction enzymes SmaI and Bg/II. This cloning yielded plasmid p2033 (Figure 18).

After digestion of p2033 with EcoRI, a 4.1 kb DNA fragment was recovered comprising the DMC1 promoter, the partial MSH3 cDNA in antisense orientation with respect to the promoter and the NOS terminator. This fragment was cloned into the EcoRI restriction site of plant transformation vector pNOS-Hyg-SCV to yield plasmid p3242 (Figure 18).

C. Construction of a combined MSH6/MSH3 antisense gene under the control of a single DMC1 promoter

A 3.1 kb fragment, encoding in antisense orientation the partial AtMSH6 and AtMSH3 sequences and the 35S terminator, was isolated from pCW186 by digestion with KpnI. The fragment was treated with Klenow enzyme to blunt both ends. It was then cloned into the SmaI site of plasmid p3243 (a pNOS-Hyg-SCV derivative, illustrated in Figure 19), which had been opened to delete the region between the SmaI sites. Clones containing the fragment in the antisense orientation with respect to the DMC1 promoter (described in A(ii) above) were identified by diagnostic digestion with BamHI. The selected construct was named p3261.

Another practical way of cloning the double antisense gene is as follows. A 1 kb DNA fragment encoding the carboxyterminal part of AtMSH6 is isolated from clone 62 described in Example 1 after digestion of clone 62 plasmid DNA with BamHI, which cleaves in the 5' polylinker region flanking the partial cDNA, and with EcoRI, which cleaves within the cDNA. The isolated fragment is treated with Klenow enzyme to blunt both its ends and is cloned into the recipient plasmid p2033 or p3242. For the purpose of

cloning, the recipient plasmid may be cleaved with either Aval or Ncol and can be blunted with Klenow enzyme to produce blunt acceptor ends for fragment cloning. This cloning yields two opposite orientations of cloned fragment DNA with respect to the DMC1 promoter. These can be identified by diagnostic digestion with restriction enzymes Scal or Xmnl in conjunction with Sacl. The selected construct contains the DMC1 promoter, the combined partial cDNAs for AtMSH3 and AtMSH6 (both cloned in antisense orientation with respect to the DMC1 promoter) and the NOS terminator. If the recipient plasmid is p2033, the combined antisense gene under control the single DMC1 promoter is recovered from the vector after EcoRI digestion and is cloned into the EcoRI restriction site of pNOS-Hyg-SCV.

D. Construction of a full-length MSH3 sense gene under control of the DMC1 promoter for overexpression of functional MSH3 protein

Overexpression of MSH3 protein was shown in human cells (Marra et al., 1998, Proc. Natl. Acad. Sci. USA 95, 8568-8573) to complex all available MSH2 protein. This leaves MSH6 protein without its partner, leading to the degradation of MSH6 protein and eventually to a mismatch repair phenotype.

This phenomenon is exploited to increase homeologous meiotic recombination in Arabidopsis as an alternative to antisense inhibition of AtMSH6. For this purpose the full-length cDNA encoding AtMSH3 is isolated from plasmid pPF66 by digestion with SmaI and is cloned into the SmaI site of the DMC1 expression cassettes described in A(i).

E. Selection of Recombination markers on homeologous chromosomes of *Arabidopsis* thaliana subspecies *Landsberg erecta* (Ler), *Columbia* (Col) and C24, respectively

E(i). Visual recombination markers in Arabidopsis th. subspecies C24:

Agrobacterium mediated transformation with a T-DNA containing a 35S-GUS gene, inactivated by insertion of a 35S-Ac transposable element (Finnegan et al., 1993, Plant Mol. Biol. 22, 625-633), had yielded a C24 line in which the T-DNA construct was integrated into chromosome 2. Genetic and molecular analysis of this line shows that the Ac transposon had excised from its T-DNA locus thereby restoring GUS activity, but had re-inserted into the chromosome at a distance of 16.4 cM, where it stayed fixed (due to disablement of Ac) within the chlorina gene. Insertional inactivation of the chlorina gene caused a bleached phenotype in those plants that were homozygous for this mutation. Because of the two linked phenotypic markers, chlorina3A:Ac and GUS, this C24 line was used in crosses to wildtype Ler for analysis of meiotic homeologous recombination on chromosome 2 in conjunction with molecular recombination markers.

35 E(ii). Visual recombination markers in Arabidopsis th. Ler:

The Ler line NW1 (obtained from NASC, Nottingham, UK) contains one recessive visual marker per chromosome. i.e. an-1 on Chr.1, py-1 on Chr.2, gl1-1 on Chr.3, cer2-1

on Chr.4, and ms1-1 on Chr.5. This line is used in crosses to wildtype C24 which expresses an MMR altering gene for analysis of meiotic homeologous recombination on chromosomes 1-5 in conjunction with molecular recombination markers listed in Table 1.

Other Ler lines from NASC have several visual markers in close proximity to each other on the same chromosome. When these lines are used for hybrid production, analysis of homeologous meiotic recombination may make use entirely of visual recombination markers. Particularly suitable for crossing to C24 wildtype that is expressing a MMR altering gene are the following Ler lines:

NW22: relative markers are dis1 - (4 cM) - ga4 - (11 cM) - th1 on chromosome 1.

NW10: relevant markers are tz-201 - (5 cM) - cer3 on chromosome 5.

10

NW134, relevant markers are ttg - (4 cM) - ga3 on chromosome 5.

NW24 (abi3-1) and NW64 (gl1-1). When present in the same plant on chromosome 3, abi3-1 and gl1-1 are 13 cM apart. Since this marker combination is not available from NASC, we have combined these markers by crossing line NW24 to line NW64. The F1 offspring were allowed to self-fertilise and to produce F2 seeds. F2 Plants which carry both markers as homozygous traits on the same chromosome can be identified firstly, by germinating F2 seeds on germination medium containing selective concentrations of abscisic acid, and subsequently, by identifying among the abscisic acid resistant plants those individuals which show the glabra phenotype.

20 E(iii) Molecular recombination markers in Col, Ler and C24:

The genome of Arabidopsis thaliana is interspersed with unique base sequences arranged as simple tandem repeats. Allelic repeats can vary in length between different Arabidopsis subspecies and when amplified by PCR yield diagnostic DNA products of different length named Simple Sequence Length Polymorphisms (SSLPs). Many SSLPs have been genetically mapped and have been assigned to unique chromosome locations on the recombinant inbred map (Bell and Ecker, 1994, Genomics 19, 137-144; Lister and Deans lines, Weeds World 4i, May 1997).

In Table 1 are listed 28 mapped and established SSLPs between Ler and Col. A number of PCR primer pairs are described herein (SEQ ID NO:42 to SEQ ID NO:97) which also yielded SSLPs between C24 and Ler (19 SSLPs) or between C24 and Col (25 SSLPs), respectively. Polymorphic SSLPs can be used as molecular markers in the analysis of homeologous recombination between genomes from these subspecies.

The PCR reactions (25 µL) were carried out over 35 cycles (15 seconds at 94°C, 30 seconds at 55°C and 30 seconds at 72°C), with 0.25 U Taq DNA polymerase and 0.6 µg genomic DNA in reaction buffer containing 2 mM MgCl₂. PCR products were separated by agarose gel electrophoresis (4% ultra high resolution agarose) and visualised by ethidiumbromide staining. The results from the PCR experiments are summarised in

Table 1, which also shows the sequence of PCR primers, primer annealing temperature (Tm), PCR product length and chromosome location of SSLP (with respect to the RI map of May 1997, Weeds World 4i).

F. Production of hybrid plants

C24 plants heterozygous for chlorina3A:Ac/GUS are crossed as male to emasculated wildtype Ler to produce Ler/C24(chlorina3A, GUS) hybrid seeds.

Due to the heterozygosity of the C24 parent, only 50 % of hybrid plants have inherited the chlorina3A:Ac/GUS locus. The remaining 50% of hybrid plants are wildtype with respect to chlorina3A:Ac/GUS. Since the mutant locus is linked to a kanamycin resistance gene (contained on the same T-DNA as GUS) mutant plants can be pre-selected by germinating hybrid seeds on germination medium containing 50 mg/L kanamycin.

Ler plants homozygous for the five chromosome markers are male sterile (ms1-1) and are crossed without emasculation to wildtype C24 to produce Ler(an-1, py-1, gl1-1, cer2-1, ms1-1)/C24 hybrid seeds.

Other Ler plants, which are male fertile, are crossed after emasculation of the female parent to produce Ler/C24 hybrid seeds.

G. Introduction of MSH3 and MSH6/3 antisense genes into Arabidopsis and analysis of meiotic homeologous recombination

(i) Transformation of hybrid plants and analysis of homeologous meiotic recombination

The plant transformation vectors comprising the antisense genes described in (B) and (C) above are introduced into Agrobacterium tumefaciens strain AGL1 (Lazo et al., 1991, Bio/Technology 9, 963-967) by electroporation. Recombinant Agrobacterium clones are selected on LB medium containing 50 mg/L rifampicin and 100 mg/L carbenicillin. Selected clones are used to infect roots of Arabidopsis hybrid plants (described in (F) above) using the root transformation protocol of Valvekens et al. (1988, PNAS 85, 5536-5540) except that shoot and root inducing media contain hygromycin (10 mg/L) instead of kanamycin.

Plants regenerated from roots of hybrid plants are genetic clones of root donating plants and therefore are again genetic hybrids of two *Arabidopsis* subspecies described in 30 (F). However, in contrast to the root donating plants, the regenerated hybrid plants also contain the introduced transgene and the co-introduced hygromycin resistance gene with the latter allowing these plants to grow on hygromycin containing culture medium.

Hygromycin resistant plants are then allowed to enter the reproductive phase and to produce gametes by meiotic divisions of microspore and megaspore mothercells. At meiosis, the *DMC*1 promoter is activated and can direct the expression of antisense genes described in (B) and (C) above, leading to decreased *MSH*6 and/or *MSH*3 gene

expression. This in turn-depletes the gamete mothercells of MSH6 and/or MSH3 protein, thus causing alteration of MMR during meiotic divisions and increasing the recombination frequency between homeologous chromosomes.

Transgenic plants are then allowed to self-fertilise and to produce seeds. These 5 seeds (F2 seeds with respect to hybrid production), and the plants derived therefrom, carry the homeologous recombination events which can be identified by using the visual and molecular recombination markers described in (E) above.

In case of homeologous recombination between chromosomes of Ler and C24(chlorina3A:Ac. GUS), the analysis concentrates on chromosome 2 by selecting plants showing the visual phenotypic marker chlorina. This marker thus serves as a reference point as it indicates that respective chromosomes 2 originate from C24. Other markers, such as GUS or molecular markers, on chromosome 2 may then be used to identify chromosomal regions which are derived from the Ler chromosome as a result of homeologous recombination. F2 plants of control transformants not expressing the antisense gene(s) can be analysed in parallel and the results can be used for comparison to homeologous recombination results obtained in antisense plants.

(ii) Transformation of C24 wildtype, hybrid plant production and analysis of homeologous meiotic recombination

Introduction of MMR altering genes into wildtype C24 is done using the root transformation protocol as described in G(i) for transformation of hybrid plants. Transformed plants are selected by resistance to either 10 mg/L hygromycin (in case of transformation with T-DNA's derived from pNOS-Hyg-SCV) or to 50 mg/L kanamycin (in case of transformation with T-DNA's derived from pBIN19).

Transgenic plants are then allowed to self-fertilise and to produce seeds (T1 seeds).

25 Segregation of the antibiotic resistance gene in the T1 population then indicates the number of transgene loci. Lines with a single transgene locus (indicated by a 3:1 ratio of resistant:sensitive plants) are selected and are bred to homozygosity. This is done by collecting selfed seeds (T2) from T1 plants and subsequent testing of at least four independent T2 seed populations for segregation of the antibiotic resistance gene. T2 populations which do not segregate the antibiotic resistance gene are assumed to be homozygous for both the resistance gene and the linked MMR altering gene.

C24 plants homozygous for the MMR altering gene are then crossed to Ler lines homozygous for recessive visual markers (see E(ii)) to produce C24/Ler hybrid plants as described in (F). These F1 hybrids are then allowed to enter the reproductive phase and to produce gametes by meiotic division of microspore and megaspore mothercells. At meiosis, the DMC 1 promoter is activated and can direct the expression of antisense or sense genes described in (B), (C) and (D) above, leading to decreased MSH6 and/or MSH3 gene expression. This in turn depletes the gamete mothercells of MSH6 and/or MSH3

protein, thus causing alteration of MMR during meiotic divisions and increasing the recombination frequency between the homeologous chromosomes of C24 and Ler. Recombination events are then scored in the F2 generation.

For recombination analysis, the hybrid plants are allowed to self-fertilise and to produce F2 seeds. F2 plants are then preselected for a first visual marker. Since this marker is recessive, its visual presence indicates homozygosity for Ler DNA at the relevant locus. Those F2 plants which show this first visual marker are then analysed for the presence or absence of a second visual marker which in the Ler parent is closely linked to the first marker. Absence of the second visual marker indicates recombination between the relevant C24 and Ler chromosomes between the first and second marker. The frequency of recombination in transgenic hybrids is compared to the recombination frequency in control hybrids not expressing the MMR altering gene.

Examples of recombination analysis are the following.

The Ler line NW22(dis1, ga4, th1) is used for crosses to transformed C24.

F2 plants are preselected first for thiamine requirement (th1) and then are further analysed for re-appearance of wildtype height (loss of ga4) and/or re-appearance of normal trichomes (loss of dis1) as a result of recombination.

The Ler line NW10(tz-201, cer3) is used for crosses to transformed C24.

F2 plants are then preselected first for thiazole requirement (tz) and then are further 20 analysed for re-appearance of normal, i.e. non-shiny stems (loss of cer3) as a result of recombination.

The Ler line NW134 (ttg, ga3) is used for crosses to transformed C24. F2 plants are first preselected for dwarfish appearance (ga3) and are then analysed for re-appearance of trichomes (loss of ttg) as a result of recombination.

Ler plants homozygous for abi3-1 and gl1-1 are used for crosses to transformed C24. F2 plants are first preselected for their ability to germinate in the presence of abscisic acid and are then analysed for re-appearance of trichomes on the leaves (loss of gl1-1) as a result of recombination.

In the case of homeologous recombination between transformed C24 and the Ler line 30 NW1 (an-1, py-1, gll-1, cer2-1, msl-1), recombination analysis is similar the one described above, except that the second marker is not a visual marker but has to be a molecular marker. This is because the Ler parent carries only one visual marker per chromosome.

		TABLE 1: S	SSLP Markers in Arabidopsis thaliana Subspecies	haliana	Subspecies		
Chromosome	RI Map Position	PCR Primer Pair	Primer Sequence	Tm (°C)	length/COL	length/LER	length/C24
_	2.3	ATEAT! F	GCCACTGCGTGAATGATATG	57.8	172	162	162
		ATEAT! R	CGAACAGCCAACATTAATTCCC	58.2			
	9.3	NGA63 F	AACCAAGGCACAGAAGCG	57.3	111	89	120
		NGA63 R	ACCCAAGTGATCGCCACC	59.6			
-	40.1	NGA248 F	TACCGAACCAAAACACAAAGG	56.1	143	129	no amplific.
		NGA248 R	TCTGTATCTCGGTGAATTCTCC	58.2			
_	81.2	NGA128 F	GGTCTGTTGATGTCGTAAGTCG	60.1	180	190	no amplific.
		NGA128 R	ATCTTGAAACCTTTAGGGAGGG	58.2			
1	81.2	NGA280 F	CTGATCTCACGGACAATAGTGC	1.09	105	85	85
		NGA280 R	GGCTCCATAAAAAGTGCACC	57.8			
1	111.4	NGA111 F	CTCCAGTTGGAAGCTAAAGGG	00	128	162	170
		NGA111 R	TGTTTTTAGGACAAATGGCG	70			
11	ca. 7.5	NGA 168 F	CCTTCACATCCAAAACCCAC	\$7.8	213	217	208
		NGA 168 R	GCACATACCCACAACCAGAA	87.8			

=	ca. 48	NGA1126L	CGCTACGCTTTTCGGTAAAG	57.8	161	199	961
		NGA1126R	GCACAGTCCAAGTCACAACC	59.9			
	62.2	NGA361L	AAAGAGATGAGAATTTGGAC	51.7	114	120	114
		NGA361R	ACATATCAATATATAAAGTAGC	49.5			
				i			
11	73	NGA168 F	TCGTCTACTGCACTGCCG	59.6	151	135	135
		NGA168 R	GAGGACATGTATAGGAGCCTCG	61.9			
=======================================	ca. 77	AthBIO2 L	TGACCTCCTTCCATGGAG	59.9	141	209	139
		AthBIO2 R	TTAACAGAAACCCAAAGCTTTC	54.5			
11	ca. 83	AthUBIQUE L	AGGCAAATGTCCATTTCATTG	54.1	146	148	148
		AthUBIQUE R	ACGACATGCAGATTTCTCC	57.8			
=	3.4	NGA172 F	AGCTGCTTCCTTATAGCGTCC	98	162	136	140
		NGA172 R	CATCCGAATGCCATTGTTC	55.4			
	12.8	NGA126 F	GAAAAAACGCTACTTTCGTGG	56.1	119	147	no amplific.
		NGA126 R	CAAGAGCAATATCAAGAGCAGC	58.2			
	17.5	NGA162 F	CATGCAATTTGCATCTGAGG	55.8	107	68	no amplific.
		NGA162 R	CTCTGTCACTCTTTTCCTCTGG	1.09			

III	81.8	NGA6 F	TGGATTTCTTCCTCTCTCAC	56.1	143	123	143
		NGA6 R	ATGGAGAGCTTACACTGATC	56.1			
2.	8.61	NGA12 F	AATGTTGTCCTCCCTCCTC	59.9	247	234	220
		NGA12 R	TGATGCTCTGAAACAAGAGC	58.2			
VI	24.1	NGA8 F	GAGGCAAATCTTTATTTCGG	56.1	154	861	961
		NGA8 R	TGGCTTTCGTTTATAAACATCC	54.5			
١٧	102	NGA1107 L	GCGAAAAACAAAAAATCCA	50.2	150	140	140
		NGA1107 R	CGACGAATCGACAGAATTAGG	58			
		•					
^	11.8	NGA225 F	GAAATCCAAATCCCAGAGAGG	58	119	189	119
		NGA225 R	TCTCCCCACTAGTTTTGTGTCC	60.1			
^	16.7	NGA249 F	TACCGTCAATTTCATCGCC	55.4	125	115	115
		NGA249 R	GGATCCCTAACTGTAAAATCCC	58.2			
^	6'61	CA72 F	AATCCCAGTAACCAAACACACA	56.3	124	110	110
		CA72 R	CCCAGTCTAACCACGACCAC	61.9			
۸	20	NGA151 F	GTTTTGGGAAGITTTGCTGG	55.8	150	120	130
		NGA151 R	CAGTCTAAAAGCGAGAGTATGATG	58.6			

>	24	NGA106 F	GTTATGGAGITTCTAGGGCACG	60.1	157	123	081
		NGA106 R	TGCCCATTTGTTCTC	55.8			
>	37.8	NGA139 F	AGAGCTACCAGATCCGATGG	59.9	174	132	132
		NGA139 R	GGTTTCGTTTCACTATCCAGG	55.8			
>	50	NGA76 F	GGAGAAATGTCACTCTCCACC	60.1	231	> 250	300
		NGA76 R	AGGCATGGGAGACATTTACG	57.8			
>	61.1	ATHSO191 L	ATHSO191 L CTCCACCAATCATG	55.8	148	156	146
		ATHSO191 R	TGATGTTGATGGAGATGGTCA	53.7			
۸	81.7	NGA129 F	TCAGGAGGAACTAAAGTGAGGG	1.09	177	179	172
		NGA129 R	CACACTGAAGATGGTCTTGAGG	60.1			

CLAIMS

- 1. An isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide functionally involved in the DNA mismatch repair system of a plant.
- 2. A DNA molecule according to claim 1 wherein said polypeptide is homologous to a mismatch repair polypeptide of a yeast or of a human.
- 3. A DNA molecule according to claim 1 wherein said polypeptide is homologous to AtMSH3 (SEQ ID NO: 19) or to AtMSH6 (SEQ ID NO: 31).
- 4. An isolated and purified polypeptide functionally involved in the DNA 10 mismatch repair system of a plant.
 - 5. A polypeptide according to claim 4 which is homologous to a mismatch repair polypeptide of a yeast or of a human.
- 6. An isolated and purified polypeptide selected from the group consisting of a polypeptide encoded by the gene *AtMSH3* (SEQ ID NO: 18), a polypeptide encoded by the gene *AtMSH6* (SEQ ID NO:30), polypeptides homologous to a polypeptide encoded by the gene *AtMSH3* (SEQ ID NO: 18) and polypeptides homologous to a polypeptide encoded by the gene *AtMSH6* (SEQ ID NO:30).
- 7. An isolated and purified DNA molecule comprising a polynucleotide sequence selected from the group consisting of (i) a sequence encoding a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence; and (ii) a sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant.
- 8. A DNA molecule according to claim 7 comprising a polynucleotide sequence encoding a polynucleotide capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence.
- 9. A DNA molecule according to claim 8 wherein said polynucleotide is capable 30 of interfering with the expression of a plant polynucleotide sequence is a sense polynucleotide, an antisense polynucleotide or a ribozyme.
 - 10. A DNA molecule according to claim 7 comprising a polynucleotide sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant.

- 11. A DNA molecule according to claim 10 wherein said polypeptide is homologous to AtMSH3 (SEQ ID NO: 19) or to AtMSH6 (SEQ ID NO: 31).
- 12. A DNA molecule according to claim 10 further comprising a regulation element capable of causing overexpression of said polypeptide in a cell of said plant.
- 5 13. A chimeric gene comprising:
- a DNA sequence selected from the group consisting of (i) a sequence encoding a polynucleotide capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence, and (ii) a sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant; and

at least one regulation element capable of functioning in a plant cell.

- 14. A chimeric gene according to claim 13 wherein said regulation element is selected from constitutive, inducible, tissue type specific and cell type specific promoters.
- 15. A chimeric gene according to claim 13 comprising a DNA sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant, wherein said regulation element is capable of causing overexpression of said polypeptide in a cell of said plant.
- 16. A chimeric gene according to claim 13 wherein said regulation element is selected from the group consisting of 35S, NOS, PR1a, AoPR1 and DMC1.
 - 17. A plasmid or vector comprising a chimeric gene according to any one of claims 13-16.
 - 18. A plant cell stably transformed, transfected or electroporated with a plasmid or vector according to claim 17.
- 25 19. A plant comprising a cell according to claim 18.
 - 20. A plant according to claim 19 selected from plants of the families Brassicaceae, Poaceae, Solanaceae, Asteraceae, Malvaceae, Fabaceae, Linaceae, Canabinaceae, Dauaceae and Cucurbitaceae.
- 21. A process for at least partially inactivating a DNA mismatch repair system of a plant cell, comprising transforming or transfecting said plant cell with a DNA molecule according to any one of claims 1-3 or 7-12 and causing said DNA sequence to express said polynucleotide or said polypeptide.
 - 22. A process for at least partially inactivating a DNA mismatch repair system of a plant cell, comprising transforming or transfecting said plant cell with a chimeric gene

according to any one of claims 13-16 and causing said DNA sequence to express said polynucleotide or said polypeptide.


- 23. A process for at least partially inactivating a DNA mismatch repair system of a plant cell, comprising transforming or transfecting said plant cell with a plasmid or vector according to claim 17 and causing said DNA sequence to express said polynucleotide or said polypeptide.
- 24. A process for increasing genetic variation in a plant comprising obtaining a hybrid plant from a first plant and a second plant, or cells thereof, said first and second plants being genetically different; altering the mismatch repair system in said hybrid plant; permitting said hybrid plant to self-fertilise and produce offspring plants; and screening said offspring plants for plants in which homeologous recombination has occurred.
- 25. A process according to claim 24 wherein a first gene is incapacitated in said first plant, a second gene is incapacitated in said second plant, and said first and second genes are incapacitated in said hybrid plant thereby altering the mismatch repair system of said hybrid plant.
 - 25. A process according to claim 25 wherein said incapacitation of the mismatch repair system of said hybrid plant is reversible.
- 26. A process according to claim 24 wherein a new genetic linkage of a desired characteristic trait or of a gene which contributes to a desired characteristic trait is 20 observable in at least one of said offspring plants.
- 27. A process for obtaining a plant having a desired characteristic, comprising altering the mismatch repair system in a plant, cell or plurality of cells of a plant which does not have said desired characteristic, permitting mutations to persist in said cells to produce mutated plant cells, deriving plants from said mutated plant cells, and screening said plants for a plant having said desired characteristic.
- 28. A process according to claim 27 wherein said step of altering the mismatch repair system comprises introducing into said hybrid plant, plant, cell or cells a chimeric gene according to claim 13 and permitting the chimeric gene to express a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence in a mismatch repair gene of the hybrid plant, plant, cell or cells, or a polypeptide capable of disrupting the DNA mismatch repair system of the hybrid plant, cell or cells.
 - 29. A process according to claim 28 comprising inactivating an MSH3 gene and/or an MSH6 gene of said plant.
- 30. A process according to claim 28 comprising inactivating an MSH3 gene and an MSH6 gene of said plant.

- 31. A process according to claim 27 comprising at least partially inactivating the mismatch repair system of said plant in a predetermined cell type or in a predetermined tissue of said plant.
- 32. A process according to claim 31 further comprising restoring mismatch repair 5 in said cell type or said tissue.
 - 33. An oligonucleotide capable of hybridising at 45°C under standard PCR conditions to a DNA molecule according to claim 1 with the proviso that said oligonucleotide is other than SEQ ID NO:1 or SEQ ID NO:2.
- 34. An oligonucleotide capable of hybridising at 45°C under standard PCR to conditions to the DNA of SEQ ID NO: 18 with the proviso that said oligonucleotide is other than SEQ ID NO:1 or SEQ ID NO:2.
 - 35. An oligonucleotide capable of hybridising at 45°C under standard PCR conditions to the DNA of SEQ ID NO:30 with the proviso that said oligonucleotide is other than SEQ ID NO:1 or SEQ ID NO:2.

	·		*	٤
				· <u>·</u>
				·
,				
		·		

74.7

Figure 1



1.11




Figure 2

Comments/References: 52= 3' side of S5 (AtMSH3) 1244bp in pUC18/Sma1




Figure 3

Comments/References: 13 = 3' side of S5 (AtMSH3) 2104bp in pUC18/Sma1

•	7	ľ
	(b
	١	4
	į	3
	ì	í
1	Ġ	ų

80	1 44 15	204 35	264 55	324 75	384 95	115	504 135	564 155	624 175	684 195	744 215	804 235	864 255	924 275
AGGA	CCC	cc6 P	CTC L	GAT D	GTT V	AAG K	GGA G	TTC F	GCA A	GCA A	GAA E	AAT	GAA E	GTT V
ACGG	GCT	CCA P	CAC	200	TAT	CTA	TTC (NAT	AAT O	GGT (CTT C	AGT P	ATT G	GTT C
TTAA	TTC F	CCG	GAC D	GTA v	GAA	GAG	TTC	CAC H	GTG 1	CAT (ACG (T	CAG A	GGT 7 G]	GAA O
rrcrc	TTC F	ACA T	TCC S	CCA	GAG E	GTG V	AGA R	GAT D	CTG	TCC	GCC	TCA (TGT (GGT (
cctaagaaagcgcgcgaaaattggcaacccaagttcgccatagccacgaccacgacctt <mark>ccatttctcttaaacgga</mark> gga	CGT R	TCA S	CTC L	AAC N	CCG P	GTG V	TAC Y	ATG M	AGA R	AAG K	AAAA K	GGT G	ე ეეეე	ACA
CCTI	TCT	TCA S	CTT L	CAA O	TCG S	CAA Q	AGG R	CAT H	AGA R	ATT I	ACC T	TTT F	TTA L	TCG S
ACGA	ATT I	GAA E	AAG K	ACT	CCC	CAG Q	TAC Y	GCT A	GTG V	GCC	TAT Y	GGT G	ACA T	ATT I
GACC	ACG T	GCC	CGT R	CAC H	GAA E	GNA E	GGT G	TAC Y	CAT H	GCA A	$_{ m L}^{ m TTG}$	GAA E	GAG E	GAA E
CCAC	CAG	GTA V	AAG K	CCT P	CTG L	$_{\rm L}^{\rm TTG}$	GTT V	AT'T I	TTC F	ACT T	GCG A	GAA E	TCG	GTT V
ATAG	CAG Q	DOO d	TCC	TCT	TTT	CCA P	GAA	GGT G	AAT N	GAA E	TCG S	GGT	AAG K	ე ე
ລວອວ	A.A.G K	AAT N	CCT	CTT	AGA R	ACA T	GTG V	TTG L	${ m TTG}$	ACT T	CTG L	GGT	GTT V	GTT V
AGTT	CAA	. ccG	TCT	AAA K	CAG	TAC Y	ATG M	GTG V	CGA R	CAG Q	GGA G	TGT C	AGA R	GTT V
CCCA	AAG	GAA	TTC	CCT	CTC	AAA	TTG L	CGC R	TTT F	AAG K	CGG R	GGT	GAG	GGT G
GCAA	ງ ວອອ :	. CAC	TCC S	AAG K	TTT. F	AGG R	GTT V	GCA	ACA	GTG V	TTC	GGT G	GAT D	GTC
ATTG	. ATG	ACT T	GTA V	AAA	AGA R	TCG S	GTG V	GCA A	CCA	GTA V	TTT F	AGT S	GTG V	AGA R
GAAA	AATT	cc6	ACT T	A CCC	CAA	TCA	GAT D	ATC I	GTG V	GGT G	CCT P	ATA I	GTT V	GTT V
2525	MAAGC	rcc S) GCC	TCA S	CAC H	TCA	CCA	GAG E	AGT	ATT	9 299	GAT D	TGT C	GAT D
AAAGC	GATTACGAATAAAGCAATT ATG	, AAA K	TCC S	GCG A	TTA	ACG T	TAC	GCG A	GCG A	AAG K	ACC T	GAG E	GTT V	TTT F
PAAG	TACC	A CCC	ATA I	GCC	. AAT	GAA	AAG K	GAC D	ACG T	TAC Y	CGG R	GCT A	TTG L	AGT S
ນ	GAT	AAA K	AAG K	9 8 9	CCC	CCC P	AGC	GAA E	ATG M	GGA G	AAC	GCG A	TTC	ATG M
	81	145 16	205 36	265 56	325 76	385 96	116	505 136	565 156	625 176	665 196	745	805 236	865 256

422 Rec'd PCT/PTO 1 0 APR 2000

925 276	TAT Y	GAA E	GAG E	TTC F	AAT N	GAT D	AAT N	TTC	ATG M	AGA R	AGT (GGA 1 G I	TTA (GAG (GCT (GTG A	ATT 1 I I	TTG	AGC S	$_{ m L}^{ m TTG}$	984 295
985 296	TCA	CCA P	GCT	GAG E	CTG L	$\mathop{\mathrm{rrg}}_{\mathrm{L}}$	CTT L	9 299	CAG Q	CCT	CTT 1	TCA (CAA (CAA 1	ACT (GAG A	AAG 1 K	TTT	TTG L	GTG V	1044 315
1045 316	GCA A	CAT M	GCT A	GGA G	CCT P	ACC T	TCA S	AAC	GTT V	CGA (GTG (GAA C	CGT C	GCC 1	TCA (CTG G	GAT 1	TGT	TTC	AGC S	1104 335
1105 336	AAT N	GGT G	AAT N	GCA	GTA V	GAT D	GAG	GTT	ATT	TCA 3	TTA 1 L (TGT G	GAA P	AAA A	ATC A	AGC G	GCA G	GGT)	AAC	TTA L	1164 355
1165 356	GAA E	GAT D	GAT D	AAA K	GAA E	ATG M	A.A.G K	CTG	GAG	GCT (GCT (GAA A	AAA G	GGA A	ATG 1	rcr 1 s	16C 1	TTG /	ACA	GTT V	1224 375
1225 376	CAT H	ACA T	ATT	ATG M	AAC	ATG M	CCA	CAT	cre .	ACT (GTT (CAA Q	GCC C	CTC C	GCC (CTA A	ACG T	TTT ?	TGC	CAT H	1284 395
1285 396	CTC L	AAA K	CAG Q	TTT F	GGA G	TTT F	GAA	AGG .	ATC (CTT 1	TAC O	CAA G	000 g	GCC T	TCA 1	TTT C F R	CGC T	TCT 1	TTG	TCA S	1344 415
1345 416	AGT S	AAC	ACA	gag E	ATG	ACT	CTC	TCA (GCC A	AAT A	ACT C	CTG C	CAN	CAG I	TTG G	GAG G E V	GTT G	GTG A	AAA K	AAT N	1404 435
1405 436	AAT N	TCA S	GAT D	GGA G	TCG	GAA E	TCT (200	TCC 1	TTA 1 L	TTC C	CAT A H	AAT A N M	ATG A	AAT O	CAC A H T	ACA C	CTT ?	ACA (T	GTA V	1464 455
1465 456	TAT Y	GCT G	TCC	AGG R	CTT	CTT	AGA (CAC 1	TGG (GTG A	AC'r C T H	CAT C H P	CCT C	CTA T	TGC G	GAT A D R	AGA A R	AAT T N	TTG 1	ATA 1	1524 475
1525 476	TCT	GCT A	CGG	CTT	GAT	GCT	GTT 7	TCT (GAG P	ATT T I S	TCT G	GCT T	TGC A	ATG G M	GGA T	TCT C. S H	CAT A H S	AGT T S S	rcr 1	TCC	1584 495
1585 496	CAG Q	CTC	AGC S	AGT	GAG	TTG (GTT (GAA (GAA O	GGT T	TCT G	GAG A E R	AGA G R	GCA A	ATT G I V	GTA T	TCA C	CCT G	GAG 1	TTT F	1644 515
1645 516	TAT Y	CTC	GTG (CTC	TCC	TCA (GTC 1	TTG #	ACA G	GCT A	ATG T M S	TCT A	AGA T	TCA T S S	rcr G s D	GAT A' D I	ATT C. I Q	CAA C	CGT G	GGA G	1704 535
1705 536	ATA	ACA T	AGA R	ATC '	TTT (CAT (CGG A	ACT G	GCT A	AAA G K	GCC A	ACA G. T E	GAG T	TTC A F	ATT G I A	GCA G	GTT A	ATG G M E	GAA G	GCT A	1764 555
1765 556	ATT	TTA (CTT (L)	6CG (000	AAG (CAA A	ATT C	CAG C Q R Fi	cgg cy R L igure	Ē. 4	66C A' 6 I (Co)	C ATA AAG CA I K Q (Continued)	AAG C. K O.	Æ	GAC TOD	TCT G	GAA A E M	ATG A M R	AGG R	1824 575

PCT/EP98/06977

1825 576	AGT	ATG M	CAA	TCT	GCA	ACT T	GTG V	CGA R	TCT	ACT T	CTT L	TTG L	AGA R	AAA K	${ m TTG}_{ m L}$	ATT	TCT	GTT V	ATT I	TCA	1884 595
1885 596	TCC	CCT	GTT V	GTG V	GTT V	GAC D	AAT N	GCC	GGA G	X AA	$_{\rm L}^{\rm CTT}$	CTC	TCT S	GCC A	CTA L	AAT N	AAG K	GAA	GCG A	GCT	1944 615
1945	GTT V	CGA R	GGT G	GAC D	$_{L}^{\mathrm{TTG}}$	CTC L	GAC D	ATA I	CTA L	ATC I	ACT T	TCC S	AGC S	GAC	CAA Q	TTT F	CCT P	GAG	CTT L	GCT A	2004
2005	GAA E	GCT A	CGC R	CAA Q	GCA A	GTT V	TTA 1.	GTC V	ATC I	AGG R	gaa E	AAG	CTG L	GAT D	TCC	TCG	ATA I	GCT	TCA S	TTT F	2064 655
2065 656	CGC R	AAG K	AAG K	CTC L	GCT A	ATT I	CGA R	AAT	$_{\rm L}^{\rm TTG}$	GAA E	TTT F	CTT L	CAA	GTG V	TCG	000 0	ATC	ACA	CAT H	TTG L	2124 675
2125	ATA I	GAG	CTG L	CCC	GTT V	GAT	TCC	AAG K	GTC V	CCT P	ATG H	AAT	TGG W	GTG V	KAAA	GTA V	AAT	AGC	ACC	A.A.G K	2184 695
2185 696	AAG K	ACT	ATT	CGA R	TAT Y	CAT H	೧೦೧ ಕ	CCA P	GAA E	ATA I	GTA V	GCT A	.; 000 0	7 TG	GAT D	GAG	CTA L	GCT A	CTA	GCA	2244 715
2245 716	ACT	GAA E	CAT H	CTT L	GCC	ATT	GTG V	AAC	CGA R	GCT A	TCG S	TGG W	GAT D	AGT S	TTC F	CTC	AAG K	AGT	TTC F	AGT	2304 735
2305	AGA R	TAC Y	TAC Y	ACA T	GAT D	TTT F	AAG K	GC'F A	GCC A	GTT V	CAA	GCT A	CTT	GCT A	GCA	CTG	GAC	TGT C	TTG L	CAC H	2364 755
2365 756	TCC S	$_{\rm L}^{\rm CTT}$	TCA	ACT T	CTA L	TCT S	AGA R	AAC	AAG K	AAC	TAT Y	GTC V	CGT	CCC	GAG E	TTT	GTG V	GAT D	GAC	TGT C	2424 775
2425	GAA E	CCA P	GTT V	GAG E	ATA I	AAC	ATA I	CAG Q	TCT	GGT G	CGT R	CAT H	CCT	GTA V	CTG L	GAG	ACT	ATA	TTA	CAA 2	2484 795
2485 796	GAT D	AAC	TTC	GTC V	CCA P	aat n	GAC	ACA	ATT I	${ m TTG}$	CAT H	GCA	GAA	999	GAA E	TAT	76C	CAA	ATT	ATC I	2544 815
2545 816	ACC T	GGA G	CCT	AAC	ATG M	GGA G	GGA G	A.A.G K	AGC S	TGC C	TAT Y	ATC	CGT	CAA	GTT	GCT	TTA L	ATT	TCC	ATA I	2604 835
2605 836	ATG M	GCT A	CAG Q	GTT V	GGT G	TCC	TTT E	GTA V	CCA	900 8	TCA S	TTC	GCC A	AAG K	CTG	CAC (GTG V	CTT	GAT (GGT G	2664 855
2665 856	GTT V	TTC	ACT	000 8	ATG M	GGT G	GCT A	TCA	GAC D	AGT S igus	~	CAG	CAT H onti	CAG CAT GGC AG Q H G R (Continued)	Æ	AGT I	ACC	TTT	CIA (GAA	2724 875

•

2784 895	2844 915	2904 935	2964 955	3024 975	3084 995	3144 1015	3204 1035	3264 1055	3324 1075	3397 1082	3458 5	3522 16
CTT GTT ATA TTA	TAT GCA ACA TTA Y A T L	TAC CCT GAA ATA Y P E I	TCG TAT CTG ACA S Y L T	TAT AAG CTT GTG Y K L V	CAG ATA CCT CCA Q I P P	GTA CGT GCA AGA V R A R	GGC GCA GAA GAA G A E E E	TCT GAA GAG GAC	AAA ATC AGA CTA 3 K I R L 1	TGA TTTAATCTTAACATTATAGCAACTGCAAGGTCTTGATCATCTGTTAGTTGCG	TCT AAT CCG S N P	TAA AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
CGT TCG C	ATT GCC I	ACG CAT T	CAT GTC T	TAC CTA T Y L Y	CTT GCC C	GCT GAG G A E V	CCG AGA G	GCT CTC TO A	GCT GGC A	GTCTTGAT	G ATG GAT M D	аааааааа 1)
TCT TCT S S	GTA GCC V	TTT GTC ?	ACA TAC (GTG ACC	GCT CAG (TTG GAA C	GAA GAA C	AAA T'IT C K F A	AAG ATT G K I P	AACTGCAA G	TAG AGAG ATG	aaaaaaaa ontinue
ACC TGT T C	GAC GGT D G	GTT CTT V L	GTT GGG V G	GAT GAT D D	AAG GTT K V	GCA AAA A K	GGA CAT G H	GAC CTG D L	GCT TGG A W	attatagci	AAA GAG NAT K E N	: GCA TAA AAAAAAAAAAAAAA A + Figure 4 (Continued)
A ATC AGA I R	C ACA CAC T H	A TGT TTG C L	A GGT TCT G S	r GAT CAT D H	r ggt ttt g F	A ATG GCT M A	A CCA GAA P E	TTT GCA	TTA AAG CAT L K H	AATCTTAAC	ACA AGA AA T R K	CCC GCA P A Figu:
TCA CAC ATA S H I	GGC ACT AGC G T S	GAA AAG AGA E K R	GGA TTC CCA G F P	GGC AGT TAT G S Y	AGG AGC TTT R S F	GCC ATT TCA A I S	ATG GGA GAA M G E	T GAC TTG D L	TTT F	T TGA TTT	AGT ATA S I	T TCT CCA S
GAA GCG TO	GGA AGA GC G R G	CTA GCA G	AGT AAC GC S N G	GAT AAA GC D K G	TGC AGC AC	CGT CGA GC	ACA CGC AT T R M	GCT C'FA GGT A L G	GCA TTC GAG A F E	TGT TCA TTT C S F	ATT	TAC ATC TTT Y I F
TTA AGT G	GAG CTT G	CAT CTC C	GAG ATC A	CAG AAG G	GGT CIT I	TGT ATA C	AGA AAT A R N T	ATT TCG G	TGG AAA G W K A	CCA ACT TO	TACTAACTT ATG TGT M C	TTG CAG TI L Q Y
2725 GAA 876 E	2785 GAT 896 D	2845 CAG 916 Q	2905 GCT 936 A	TTG L	CGT R	TCA	GAG E	TCT	CCT	AAA K		GTG V
0.00	27	28	9.9	2965 956	3025 976	3085 996	3145 1016	3205 1036	3265 1056	3325 1076	3398	3459 6

Agree of the

2S8 AP2 AP1 clone 62_ 1382bp 637 S823 638 Xmn1 S82 S83 3330bp Figure 6 clone 43 – 2182bp API AP2 158 581

SURSTITUTE SHEET (RIII E 28)




Figure 7

Comments/References: 43= 5' side of S8 (AtMSH6) 2182 bp in pUC18/Sma1




Figure 8

Comments/References: 62= 3' side of S8 (AtMSH6) 1379bp in pUC18/Sma1

with the state of the state of

1	AAA	aaaagttgagccctgaggagtatcgtttccgccatttctacgacgcaaggcgaaatttttggcgccaafctttcccccc	BAGCC	CTGA	GGAG	STATO	GTTT	ວອວວຸ	CATT	TCTA	CGAC	GCAA	ວວອອ	AAAA	TTTT	TGGC	GCCA	ATCT	TTCC	၁၁၁၁	80
81	TTT	TTTCGAATTCTCTCAGCTCAAAACATCGTTTCTCTCTCACTCTCTCT	rrcrc	TCAG	CTC	AAAAC	ATCG	TTTC	TCTC	TCAC	TCTC	TCTC	ACAA	TTCC	AAAA	A ATG M	G CAG	ဂ င ၆င R	c cag	ပ	153 4
154	AGA R	TCG S	ATT	$_{ m L}^{ m TTG}$	TCT S	TTC	TTC	CAA	AAA K	CCC	ACC T	909 8	GCG	ACT	ACG	AAG	GGT G	TTG L	GTT V	TCC S	213 24
214 25	9 299	GAT D	GCT A	GCT A	AGC	ე ე	9 9	ე ეეე	255	AGC S	GGA G	GGA	CCA P	CGA R	TTT F	AAT	GTG	AAG R	GAA	9 9	273
274 45	GAT D	GCT A	AAA K	၁၅	GAC	GCT	TCT	GTA V	CGT	TTT F	GCT	GTT V	TCG	AAA K	TCT	GTC	GAT D	GAG	GTT V	AGA R	333 64
334 65	GGA G	ACG T	GAT D	ACT	CCA P	CC6	GAG E	AAG K	GTT	CCG	CGT R	CGT	GTC V	$\mathop{\mathtt{CTG}}_L$	CCG P	TCT	GGA	TTT	AAG K	CCG P	393 84
394 85	GCT A	GAA E	TCC	GCC	GST G	GAT D	GCT	TCG S	TCC	CTG L	TTC F	TCC	AAT	AT'T I	ATG M	CAT	AAG K	TTT	GTA V	AAA K	453 104
454 105	GTC V	GAT D	GAT D	CGA R	GAT D	TGT	TCT S	GGA G	GAG	AGG	AGC S	CGA R	GAA	GAT	GTT V	GTT (SCC B	CTG	AAT	GAT D	513 124
514 125	TCA	TCT	CTA L	TGT C	ATG M	AAG K	GCT A	AA'I'	GAT (GT'T V	AT'r I	CCT (CAA	ř.ř.ř	CGT	rcc s	AAT	AAT	GGT G	AAA K	573 144
574 145	ACT T	CAA Q	GAA E	AGA R	AAC	CAT H	GCT	TTT	AGT '	TTC	AGT	666 J	AGA (GCT (GAA E	CTT	AGA '	TCA (GTA	GAA E	633 164
634 165	GAT D	ATA I	GGA G	GTA V	GAT D	၁၅၅	GAT	GTT V	CCT (GGT	CCA P	GAA)	ACA (CCA (999	ATG (CGT (CCA (CGT	GCT	693 184
694 185	TCT	CGC R	${ t TTG}$	AAG K	CGA R	GTT V	CTG	GAG (GAT (GAA	ATG	ACT 7	TTT /	AAG (GAG (GAT 1	AAG (GTT (CCT	GTA V	753 204
75 4 205	$\mathop{\mathrm{TTG}}_{L}$	GAC D	TCT	AAC	AAA K	AGG R	CTG	AAA I	ATG (CTC (CAG (GAT (500	GTT 7	TGT (GGA (GAG 1	AAG 7	AAA	GAA E	813 224
814 225	GTA V	AAC	GAA	GGA	ACC	AAA K	TTT (GAA '	TGG &	CTT (GAG	TCT 7	TCT (CGA)	ATC i	AGG OR	GAT (GCC A	AAT	AGA R	873 244
874 245	AGA R	CGT R	CCT	GAT	GAT D	CCC	CTT .	TAC (GAT A	AGA 1	AAG 1	ACC 1	TTA (CAC A	ATA (CCA (CCT C	GAT (GTT	TTC	933 264

ATT 993 I 284	GGT 1053 G 304	GGT 1113 G 324	GTT 1173 V 344	ATA 1233 I 364	GGG 1293 G 384	TCA 1353 S 404	ATC 1413 I 424	GAA 1473 E 444	ACG 1533 T 464	GAT 1593 D 484	TCA 1653 S 504	GAG 1713 E 524	TTT 1773 F 544	GAG 1833
GAC .	TTA L	GTT V	AAA K	ACT T	ATC I	$ au_{\mathrm{C}}$	TCC	AAG K	TAT Y	ACA T	gaa E	GGA G	ATT I	CTT
ATG G M	GAA E	CAG Q	TAT Y	AAT N	AAC	AAG K	9 9	CCA P	AAA K	GAT	TCT	CTT (GAT D	AAT (
rat A Y	GCG A	AGA R	GGA G	GCT A	GGA G	CAA	GTT V	TCT	AGG R	999	TCT	GCT () 999	GTA A
SAA 1	GAT D	TGC C	CGT R	GGT G	GAG E	CTA L	TGG ₩	GTT V	CTA	ATG	GGT	AGT (CAT (ATG (
AGT C	CTA L	AAA K	GCT A	AGA R	AGC	GAG E	TTT F	CAG Q	GCT A	GTA V	AAA K	CTT	AAG (ACG 1
AAG 1 K	GAG E	GGA G	$_{\rm L}^{\rm TTA}$	GCC	GCA	ATG M	AGG R	ATG M	AAG K	CAN	TTT F	GCC	CTT	CAG
GTT /	TAT Y	GTG V	CTA L	A.A.A	ACA T	A A A	TTG L	$_{\rm L}^{\rm TTG}$	CAA	CCA P	TAC Y	GTT	GTA V	255
AGT (CTG L	GGT G	AAG K	GCA A	TCA S	ATC	GCC	TTA L	GCA A	GTA V	GGA G	GAT D	GAT D	GAT
TGG ₹	GAG E	AGT S	CAA Q	CAA	CCA P	GAG E	GCT A	gcc A	GAA E	CCA P	AAC N	TGT	GAA E	ATT
TAT '	TAT Y	ATG M	GTG V	GAC D	ACT T	AAA K	TGT C	GGA G	AGA R	GCT	TCT S	GAA E	CTA	AGA
CAA	TTT F	ACC T	GCA A	TCT S	TTA L	ATA I	GAC D	CTT L	TCA S	TTG L	GAA	AAT N	AAG K	CTC
AAG	AAA K	ATG M	GAG	ACA T	GTA V	GCT A	GTT V	GCT A	CTA L	CAG Q	ATA I	CTA L	CTA L	TGT
CAS	ე ე	A.A.G K	GAT D	gaa E	CAG Q	$_{\rm L}^{\rm CTT}$	TTT F	GCT A	9 999	GTA V	ATA I	GGT G	AGG R	GGT
rca s	GTG V	TGG W	ATA I	CTA L	GTT V	CTT	GCT A	TGT C	AAA K	GCG A	AAT N	GAT	TCT	AGG
GCA '	AAA K	GAC	9 999	CAG Q	CTA	CAT H	TTT F	TCA S	AGT S	ACG T	AGA R	GTT V	CTG L	TAC
TCT (TTT F	CTT L	AGT S	GAG E	AAG K	GTC V	GGA G	GCA A	GAC D	TCT	GTT V	GCT A	CAT H	GTT
ATG '	TTC F	GAG E	GAA E	ATC I	AGG R	GCC	TAT Y	GAT D	TAT Y	999 9	GGA G	TGT C	AAT	CAA
AAA K	CTT L	AAG K	TCT S	CGA R	CCA P	GAT D	GTG V	GAT D	TTA L	ACA T	GCT A	AAC	ATT I	TAC
AAG K	GTG V	CAC H	ATC	GGA G	ATT I	CCT P	ACT T	AGC S	GTG V	$ ext{TTG}$	GCT GCT (TGG W	CTA	CCA
934 265	996 285	1054 305	11114 325	1174 345	1234 365	1294 385	1354 405	1414	1474	1534 465	1594 485	1654 505	1714 525	1774

•

*1

1834 565	ATA I	TTT F	AAC	AAT N	AGC S	TGT C	GAT D	GGT G	GGT G	CCT	TCA S	999	ACC T	TTG L	TAC Y	AAA K	TAT (Y	CTT (GAT D	AAC N	1893 584
1894 585	TGT C	GTT V	AGT S	CCA P	ACT	GGT G	A.A.G K	CGA R	CTC L	T'ſA L	AGG R	AAT' N	TGG W	ATC I	76C C	CAT (CCA (CTC 1	AAA K	GAT D	1953 604
1954 605	GTA V	GAA E	AGC S	ATC I	AAT N	AAA K	CGG R	$_{\rm L}^{\rm CTT}$	GAT D	GTA V	GTT V	GAA .	GAA E	TTC	ACG T	GCA 1	AAC N	TCA (GAA	AGT S	2013 624
2014 625	ATG M	CAA O	ATC I	ACT T	9 299	CAG	TAT Y	CTC L	CAC H	AAA	CTT	CCA O	GAC D	TTA	GAA	AGA (CTG (CTC (GGA (cGC R	2073 644
2074 645	ATC	AAG K	TCT	AGC S	GTT V	CGA R	TCA S	TCA S	GCC	TCT	GTG V	TTG (CC:T P	GCT	CTT	CTG (2 555	AAA I	AAA (GTG V	2133 664
2134 665	CTG	A A A	CAA	CGA R	GTT V	A.A.A K	GCA A	TTT F	0 0 0	CAA	ATT I	GTG ,	AAA K	6 6	TTC	AGA A	AGT (GGA 7	ATT (GAT D	2193 684
2194 685	CTG L	TTG L	TTG L	GCT A	CTA L	CAG Q	AAG K	GAA E	TCA	AAT	ATG	ATG /	AGT S	TTG . L	CTT	TAT A	ANA C	CTC 1 L (TGT 1	A.A.A K	2253 704
2254 705	$_{\rm L}^{\rm CTT}$	CCT P	ATA I	TTA L	GTA V	GGA G	AAA K	AGC S		CTA L	GAG	TTA	TTT (F	CTT	TCT	CAA 1	TTC C	GAA (GCA (GCC A	2313 724
2314 725	ATA I	GAT D	AGC S	GAC D	TTT F	CCA P	AAT N	TAT Y	CAG	AAC I	CAA (GAT (GTG /	ACA (GAT (GAA P	AAC G	GCT G	GAA I	ACT T	2373 744
2374 745	CTC	ACA T	ATA I	CTT L	ATC 1	GAA	C.F.T. L	TTT F	ATC (GNA	AGA (GCA A	ACT (CANA	TGG .	TCT G	GAG G E V	GTC P	ATT C	CAC H	2433 764
2434 765	ACC	ATA I	AGC S	TGC C	CTA L	GAT	GTC	CTG	AGA '	TCT	TTT (F	GCA A	ATC (GCA (GCA 1	AGT C	CTC T	TCT G	GCT G	GGA G	2493 784
2494 785	AGC S	ATG M	GCC A	AGG R	CCT	GTT	ATT	TTT	CCC (GAA 3	TCA (GAA (GCT A	ACA (T	GAT (CAG A	AAT C	CAG A	AAA A K	ACA T	2553 804
2554 805	AAA K	999	CCA P	ATA	CTT	AAA K	ATC (CAA (GGA (CTA 1	TGG (CAT C	CCA 1	TTT (F	GCA (GTT G	GCA G A A	GCC G	GAT G	GGT G	2613 824
2614 825	CAA	${ m TTG}_{ m L}$	CCT	GTT	CCG	AAT	GAT D	ATA (I	CTC C	CTT (9 0 0	GAG G	GCT P	AGA P	AGA A	AGC A	AGT G S G	300 A	AGC A	ATT I	2673 844
2674 845	CAT H	CCT	CGG R	TCA	TTG L	TTA L	CTG 1	ACG (GGA (CCA P P P Figu	AAC A N N	ATG G M G	G GGC G G C	GGA P G F nued)	AAA 3	CA A	ACT C	CTT C	CTT C	cgT R	2733 864

2734 865	GCA	ACA T	rgr c	CTG L	gcc A	GTT V	ATC I	TTT F	GCC	CAA O	CTT L	ე ე	TGC C	TAC Y	GTG V	CCG P	r_{GT}	GAG E	TCT	TGC C	2793 884
2794 885	GAA	ATC	TCC	CTC L	GTG V	GAT D	ACT T	ATC	TTC	ACA	AGG R	$_{\rm L}^{\rm CTT}$	ე ე	GCA A	TCT	GAT D	AGA R	ATC	ATG M	ACA T	2853 904
2854 905	GGA G	GAG E	AGT S	ACC T	TTT F	${\overset{\text{TTG}}{\text{L}}}$	GTA V	GAA E	TGC	ACT T	GAG E	ACA T	GCG A	rca s	GTT V	CTT	CAG Q	AAT	GCA A	ACT T	2913 924
2914 925	CAG D	GAT D	TCA S	CTA L	GTA V	ATC I	$_{\rm L}^{\rm CTT}$	GAC	GA.A E	CTG L	ე ეეე	AGA R	GGA G	ACT	AGT S	ACT T	TTC	GAT	GGA G	TAC Y	2973 944
2974 945	9 8	ATT	GCA	TAC Y	TCG	GTT V	TTT F	CGT R	CAC	CTG L	GTA V	GAG E	AAA K	GTT	CAA	TGT	CGG	ATG	CTC	TTT E	3033 964
3034 965	GCA A	ACA T	CAT H	TAC Y	CAC H	CCT P	CTC	ACC T	AAG K	GAA E	TTC	GCG	TCT S	CAC	CCA P	CGT R	GTC	ACC	TCG	AAA K	3093 984
3094 985	CAC	ATG	GCT	TGC C	GCA	TTC F	AAA K	TCA S	AGA R	TCT	GAT	TAT Y	CAA	CCA P	CGT R	GGT	TGT	GAT	CAA	GAC D	3153 1004
3154 1005	CTA L	GTG V	TTC	$\mathop{\rm TTG}_{\rm L}$	TAC Y	CGT R	TTA L	ACC	GAG E	GGA G	GCT	TGT	CCT	GAG	AGC S	TAC (GGA	CTT (CAA	GTG V	3213 1024
3214 1025	GCA A	CTC L	ATG M	GCT A	GGA G	ATA I	CCA P	AAC	CAA O	GTG V	GT'I V	GAA	ACA (GCA	TCA	GGT (GCT (GCT (CAA	GCC A	3273 1044
3274 1045	ATG M	AAG K	AGA R	TCA S	ATT I	ე ეეე	GGA E	AAC	7"rC F	AAG	TCA	AGT' I	CAC (E	CTA	AGA R	TCT (GAG :	TTC	TCA	AGT	3333 1064
3334 1065	CTG L	CAT H	GAA E	GAC	TGG W	CTC L	AAG K	TCA S	TTG L	GTG	GGT	ATT	rcr (CGA (GTC (GCC (CAC 1	AAC I	AAT N	GCC A	3393 1084
3394 1085	CCC P	ATT I	299	GAA	GAT	GAC D	TAC Y	GAC	ACT	TTG	TTT '	TGC	TTA 1	TGG (CAT (GAG A	ATC A	AAA 7	rcc 3	TCT S	3453 1104
3454 1105	TAC Y	TGT	GTT V	CCC	AAA K	TAA *	ATG	GCT A	ATG	ACA '	TAA (CACT	ATCT(3AAG(CTCG'	cactatctgaagctcgttaagtcttttgcctctct	GTCT	rtrg(CCTC	rcr .	3521 5
3522	G ATG	G TTT	T ATT I	T CCT	T CTT L	T AAA K	a aaa K	A TGC	C TTA L	A TAT Y	T ATC I	C AAA K	A AAA K	A TTG	G TTT F	r cct P	r cga R	A TTA L	A AAA K	Gr.	3579 19
3580 20	AAA K	AAA K	AAA K	A.A.A	AAA K	AAA K	A A A	AAA K	AAA K	Figure		<u>ပ</u>) 6	(Continued)	(pen						-	3606 28

\$
_
ຍ
=
\supset
20
==
_

Mark Cares		LASSABARANTANDE CONTROL OF THE TAX OF THE TA
ACE SECTION SE	***	Perverripgor Gearglocate Dalmen
ATMENS - BL	163	Vedigyddyngertegerregyleden yn kurykruk van karken van kun kolos Caryes korkorkorkorkar en en san en en en en e Deretarkryskeres san san san karak en
ACKBES - BA MAKE: Year	3 2 2	- arsokovnavkskumpetvestvovnovnovnovnovnovnum sedvokerovena sa sabarostera svenska sa seda kang. Beargir Dru - pre kovnavkskumbetvestvovnovnovnovnovnovnovnovnovnovnovnovnovn
19 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	305	ээг үндээдлэндийн тойун цамийн баган индаруулар түүндүй байдар майдардан бан байдан байдан байдан байдан байдан 117 жыз одаан байдан байдадаган индаруулуу тойун байдар майдайдайдан байдан байдан байдайдайдайдагайдагайдага
14 - 多耳病医丛皮	222	KYTLTOSTAVQLAPVPOVKODTDAAOVAKIRESAGYZKUSSERATAVULKKU DŽALŠKILGARETARETARAKIRD VLGRODIPPYOVYKULIG 100.50 : prbaprasiparvaurry Duditare išiliskuls - trijožpykulkatipiukažos ižkirio duvykumentrakire topovksukavligi. Ilu
ACEMBE - BA	223	HERE BESCHOOF FOR DELEGENEET BESCHERFEREN FOR STEELEN STEELEN SESHOLF OF THE FORESTEELEN SESHOLF OF THE STEELEN SESHOLF OF THE STEELE SESHOLF OF THE STEELEN SESHOLF OF THE STEELE STEELEN SESHOLF OF THE STEELEN SESHOLF OF THE STEE
10年4日末 中京町東 10年 - 18京東京10年	711	elgoslemmolkodvert 18sppellusky Lkoskykago 1ykgpreshioluskork
ACKUMO MO-MI Man 6. Year	725	1 dede pakkanteren 180 m dokko-kanteren 181 paken 188 kokanteren 186 meteren 187 meteren 188 meteren 188 meter Attavesammen meteren 180 metaken 188 meteren 188 me
ATMENTS - BA	9 0 0	garaldriggerleggerleverspringledarrssusihPRsleggerleggerleggerleggerleggerleggerleggerleggerleggerleggerlegger glaussyndergerleggerleggerleggerleggerleggerussussusihPRsleggerlegkangskom Grypegsgerlesgerleggerleggerlegger
ACKMENT - BL	1016	905 GESTEPVECTEVASVEDMATODESEVILDELGROTET FOGYATAY SVFRR VOCKKLEVET RANGERSERA GERENAL STEPVES STEPVES SOLUTIVET STEPVES SOLUTIVET STEPVES SOLUTIVE SVET STEPVES SOLUTIVE SVET STEPVES SOLUTIVE SVET STEPVES SOLUTIVE SVET SVET SVET SVET SVET SVET SVET SV
Aで放送される 開放の「どのものな	1010	1010 bltsocheppe, chile bos novvet Deum. 1111 kriegosechebken om vet Deum.
ACMBEG-BI	1 103	ST TDDL9

TTTTTTGGTTGCTAACAATAAAGGTATACGGTTTTATGTCATCAATATAA	50
CTATATATAAAAGAAATGAAAGATATATATTGTTTTTTCATTTATCAAAC	100
AAAACAACAAGACTTTTTTTTTTTTTTTTTTTTTTTTTT	150
GATAAACGACATCGTTTAATCATTTCCCAATTTTACCCCTAAGTTTAACA	200
CCTAGAACCTTCTCCATCTTCGCAAGCACAGCCTGATTAGGAACAGCTTT	250
ACCATTCTCATATTCCTGAACTACCTGAGTCCTCTCATTGATCTGTTTCG	300
CCAAATCCGCTTGTGACATCTTCTTCTCCAATCTCGCTTTCTGTATCATC	350
AACCTCACCTCTGCTTTCACACGATCCATCGCCGCAGGCTCTGTTTCTTC	400
TTCCAGCTTCTTCGTGTTAATCACCGGAACCGCCGTAGATTTCCCCTTTT	450
TGTTCGAACCGGCATCGAATTTCTTAACCGTTTGAACCGCGACACCGTTT	500
CTCAGAGCTGCGTTAACCGCTTTCGGATCGCGTAGGTCTTGGCTCTTTTG	550
TTTTGATTTGTGGAGAACTACTGGTTCCCAGTCTTGTGTTACTGCTCCTG	600
GGTATCTGCTCGGCATCGTCGATGAATTGAGAGAAAGGAACAACGCGAAA	650
ATTTTATTAATCTGAGTTTTGAAATTGAGAAACGATGAAGATGAAGAATG	700
TTGTTGAGAGGATTGTGATATTTATATATACGAAGATTGGTTTCTGGAGA	750
ATTCGATCATCTTTTCTCCATTTTCGTCTCTGGAACGTTCTTAGAGATG	800
ATTGACGACGTGTCATTATCTGATTTGCAGTTAACCAATGCTTTTTGGGT	850
TGGATTCGTGGTACACCATATTATCCGATTTGGCTCAATGGTTTTATATA	900
AATTTGGTTTTCGGTTCGGTTATGAGTTATCATTAAAATTAAGCTAACCA	950
AAAATTTTCGTAAAATTTATTTCGGTTTCAATTCGGATCCCTTACTTCCA	1000
GAACCGAATTATTCGAAACCGGGGTTAGCCGAACCGAATACCAATGCCTG	1050
ATTGACTCGTTGGCTAGAAAGATCCAACGGTATACAATAATAGAACATAA	1100
ATCGGACGGTCATCAAAGCCTCAAAGAGTGAACAGTCAACAAAAAAGTT	1150
GAGCCCTGAGGAGTATCGTTTCCGCCATTTCTACGACGCAAGGCGAAAAT	1200
TTTTGGCGCCAATCTTTCCCCCCTTTCGAATTCTCTCAGCTCAAAACATC	1250
GTTTCTCTCTCACTCTCTCACAATTCCAAAAAATGCAGCGCCAGAGAT	1300
CGATTTTGTCTTTCCAAAAACCCACGGCGGCGACTACGAAGGGTTTG	1350
GTTTCCGGCGATGCTGCTAGCGGCGGGGGGGGGGGGGGG	1400
AATGTGAAGGAAGGGGATGCTAAAGGCGACGCTTCTGTACGTTTTGCTGT	1450
TTCGAAATCTGTCGATGAGGTTAGAGGAACGGATACTCCACCGGAGAAGG	1500
TTCCGCGTCGTGTCCTGCCGTCTGGATTTAAGCCGGCTGAATCCGCCGGT	1550
GATGCTTCGTCCCTGTTCTCCAATATTATGCATAAGTTTGTAAAAGTCGA	1600
TGATCGAGATTGTTCTGGAGAGAGGTACTAATCTTCGATTCTCTTAATTT	1650
TGTTATCTTTAGCTGGAAGAAGAAGATTCGTGTAATTTGTTGTATTCGTT	1700
GGAGAGATTCTGATTACTGCATTGGATCGTTGTTTACAAATTTTCAGGAG	1750
CCGAGAAGATGTTGTTCCGCTGAATGATTCATCTCTATGTATG	1800
ATGATGTTATTCCTCAATTTCGTTCCAATAATGGTAAAACTCAAGAAAGA	1850
AACCATGCTTTTAGTTTCAGTGGGAGAGCTGAACTTAGATCAGTAGAAGA	1900
TATAGGAGTAGATGGCGATGTTCCTGGTCCAGAAACACCAGGGATGCGTC	1950
CACGTGCTTCTCGCTTGAAGCGAGTTCTGGAGGATGAAATGACTTTTAAG	2000
GAGGATAAGGTTCCTGTATTGGACTCTAACAAAAGGCTGAAAATGCTCCA	2050
GGATCCGGTTTGTGGAGAGAAGAAGAAGTAAACGAAGGAACCAAATTTG	2100
AATGGCTTGAGTCTTCTCGAATCAGGGATGCCAATAGAAGACGTCCTGAT	2150
GATCCCCTTTACGATAGAAAGACCTTACACATACCACCTGATGTTTTCAA	2200

GAAAATGTCTGCATCACAAAAGCAATATTGGAGTGTTAAGAGTGAATATA	2250
TGGACATTGTGCTTTTCTTTAAAGTGGTTAGTAACTATTAATCTAGTGTT	2300
CAATCCATTTCCTCAATGTGATTTGTTCACTTACATCTGTTTACGTTATG	2350
CTCTTCTCAGGGGAAATTTTATGAGCTGTATGAGCTAGATGCGGAATTAG	2400
GTCACAAGGAGCTTGACTGGAAGATGACCATGAGTGGTGTGGGAAAATGC	2450
AGACAGGTAAATTAGTTGAAACAACTGGCCTGCTTGAATTATTGTGTCTA	2500
TAAATTTTGACACCACCTTTTGTTTCAGGTTGGTATCTCTGAAAGTGGGA	2550
TAGATGAGGCAGTGCAAAAGCTATTAGCTCGTGGGTAAGGGAACCATCAT	2600
ACTTTATGGAATTCGTTTACTGCTACTTCGGCTAGGATTTAAGAAATGGA	2650
AATCACTTCAAGCATCATTAGTTAGGATCCTGAGAACTCAGGATGTTTTC	2700
TTATTCGTTATATAATAAGTCTTTTCATCAAGGAGTAACAAACA	2750
GCACAATATTTGTGTGCTCACTGGCAAGGCATATATACCCAGCTAACCTT	2800
TGCTAGTTCACTGTAGTAACAGTTACGGATAATATATGTTTACTTGTATG	2850
TGGTACCCTCATTTTGTCTCTCATGGAGGCTTTGAAGCCTTGTGTTGAAA	2900
CTGGATAGTTACATATGCTTCCAACAGAAACTAGCATGCAGATTCATATG	2950
CTTTCCTATTCTACTAATTATGTATTGACACACTCGTTGTTTCTTTTGAA	3000
AGATATAAAGTTGGACGAATCGAGCAGCTAGAAACATCTGACCAAGCAAA	3050
AGCCAGAGGTGCTAATACTGTAAGTTTTCTTGGATAGGTCAAGGAGAGTG	3100
TTGCAGACTGTTTTTGATCATTTCTTTTTTTTGTACATTACTTTCATGCTG	3150
TAATTAACTCAATGGCTATTCTGGTCTGATTATCAGATAATTCCAAGGAA	3200
GCTAGTTCAGGTATTAACTCCATCAACAGCAAGCGAGGGAAACATCGGGC	3250
CTGATGCCGTCCATCTTCTTGCTATAAAAGAGGTTTGTTATTTACTTATT	3300
TATCTTATCATGTTCAGTTCATCCAAGTCCTGAAAAATTACACTCTTCTT	3350
TACCAATCTTCCATCAAGCTGTGTAAAGGATTTGGAATTAGAAAATCATT	3400
ATTTGATGCTTTGTTTTATATGCAAGAGGTTCCCTTGAAAAGATCTGTTT	3450
AAGATTCTTTGCACTTGAAAAATTCAATCTTTTTAAGTGAATCCCCTACT	3500
TTCTTACAATGATCATAGTCTGCAATTGCATGTCAAGTAATATCATTCCT	3550
TGTTACTGCATCCCCCTCTTTCTTAATGACCATTGTCTATGTTGTGTTTG	3600
TCTCGTGTGCTGGAGAAAATGATAGCTGATCCAAGCTGTACATTATCATG	3650
ATTAAGTAGCTGCTCAGGAATTGCCTTTGGTTACATTGCCTAATGGTTTG	3700
ATGTCAATTTTCTTCTGAATCTTTATTTTAGATCAAAATGGAGCTACAA	3750
AAGTGTTCAACTGTGTATGGATTTGCTTTTGTTGACTGTGCTGCCTTGAG	3800
GTTTTGGGTTGGGTCCATCAGCGATGATGCATCATGTGCTGCTCTTGGAG	3850
CGTTATTGATGCAGGTAAGCAAGTGTATTCTGTATCTTATGTGTACCATG	3900
TGACTTCCTGTGCATATATTTGGGTTGCAGGAACTAATTCTGAATCACCA	3950
TTTGGTATGTTTTTTCCAGGTTTCTCCAAAGGAAGTGTTATATGACAGTA	4000
AAGGTAAACTGCTTGTATCGCCAGTTGTTTTGTTAAACAGAATTTAAGGT	4050
AAATGACACTGGTTAATTTAAAGTGCATACATGTTGAAATATTGCAGGGC	4100
TATCAAGAGAAGCACAAAAGGCTCTAAGGAAATATACGTTGACAGGTACC	4150
ATTTCAGTAGGCAAGCTAACTGACAATTTAACCGCTCACCGAATGATAGG	4200
TCTCTTAAACATTGCTAATGTAGATGATGTTTATGTTTCAATCTAATAGG	4250
GTCTACGGCGGTACAGTTGGCTCCAGTACCACAAGTAATGGGGGGATACAG	4300
ATGCTGCTGGAGTTAGAAATATAATAGAATCTAACGGATACTTTAAAGGT	4350
TCTTCTGAATCATGGAACTGTGCTGTTGATGGTCTAAATGAATG	4400

Figure 11 (Continued)

_

10	-				1		
	•		٠.	:	1.	 •	
	V	VO	99/1	949	2		

·		
TGCCCTTAGTGCTCTTGGAGAGCTAATTAATCATCTGTCTAGGCTAAAGG	445	0
TGTGTTGGCTTGTTTAGTTTTTGCTTTTCACAAATTAAGCAAAGGAACTT	450	0
TTCATAACTTACAGTTTCTATCTACTTGCAGCTAGAAGATGTACTTAAGC	455	0
ATGGGGATATTTTTCCATACCAAGTTTACAGGGGTTGTCTCAGAATTGAT	460	0
GGCCAGACGATGGTAAATCTTGAGATATTTAACAATAGCTGTGATGGTGG	4 6 5	0
TCCTTCAGGCAAGTGCATATTTCTTTTTTGATAACTTCAACTAGAGGGCA	470	0
GACATAGAAGGAAAAATTCTAATACTTCGTACGGATCTCCAGTAAGTA	475	o
AGCCGATTTTTGTTTACCTATGTAGGGACCTTGTACAAATATCTTGATAA	480	0
CTGTGTTAGTCCAACTGGTAAGCGACTCTTAAGGAATTGGATCTGCCATC	485	0
CACTCAAAGATGTAGAAAGCATCAATAAACGGCTTGATGTAGTTGAAGAA	490	0
TTCACGGCAAACTCAGAAAGTATGCAAATCACTGGCCAGTATCTCCACAA	495	0
ACTTCCAGACTTAGAAAGACTGCTCGGACGCATCAAGTCTAGCGTTCGAT	500	0
CATCAGCCTCTGTGTTGCCTGCTCTTCTGGGGAAAAAAGTGCTGAAACAA	505	0
CGAGTAAGTATCAATCACAAGTTTTCTGAGTAATGCCTTCCATGAGTAGT	510	0
ATAGGACTAAAACATTACGGGTCTAGCTAAAGACTGTTCTCCTTCTTTTG	515	0
CAATGTCTGGTTATTCATTACATTTCTCTTAACTTATTGCATTGCAGGTT	520	0
AAAGCATTTGGGCAAATTGTGAAAGGGTTCAGAAGTGGAATTGATCTGTT	525	0
GTTGGCTCTACAGAAGGAATCAAATATGATGAGTTTGCTTTATAAACTCT	530	_
GTAAACTTCCTATATTAGTAGGAAAAAGCGGGCTAGAGTTATTTCTTTC	535	
CAATTCGAAGCAGCCATAGATAGCGACTTTCCAAATTATCAGGTGCCCAT	540	0
CTATCTTCATACTTTACAACAAAATGTCTGTCACTACTCAAAGCAATGC	545	0
ATATGGCTTAGATCTCAACTCACCCCGAGGATCCTAAAGGGATTTGCT	550	0
TTTTATTCCTAATGTTTTTGGATGGTTTGATTTATTTCTAACTTGAACTT	555	0
ATTAATCTTGTACCAGAACCAAGATGTGACAGATGAAAACGCTGAAACTC	560	0
TCACAATACTTATCGAACTTTTTATCGAAAGAGCAACTCAATGGTCTGAG	\$65	0
GTCATTCACACCATAAGCTGCCTAGATGTCCTGAGATCTTTTGCAATCGC	570	0
AGCAAGTCTCTCTGCTGGAAGCATGGCCAGGCCTGTTATTTTTCCCGAAT	575	0
CAGAAGCTACAGATCAGAATCAGAAAACAAAAGGGCCAATACTTAAAATC	580	0
CAAGGACTATGGCATCCATTTGCAGTTGCAGCCGATGGTCAATTGCCTGT	585	0
TCCGAATGATATACTCCTTGGCGAGGCTAGAAGAAGCAGTGGCAGCATTC	590	0
ATCCTCGGTCATTGTTACTGACGGGACCAAACATGGGCGGAAAATCAACT	595	0
CTTCTTCGTGCAACATGTCTGGCCGTTATCTTTGCCCAAGTTTGTATACT	600	0
CGTTAGATAATTACTCTATTCTTTGCAATCAGTTCTTCAACATGAATAAT	605	0
AAATTCTGTTTTCTGTCTGCAGCTTGGCTGCTACGTGCCGTGTGAGTCTT	610	0
GCGAAATCTCCCTCGTGGATACTATCTTCACAAGGCTTGGCGCATCTGAT	615	0
AGAATCATGACAGGAGAGAGTAAGTTTTGTTCTCAAAATACCAATTCCTC	620	0
GAACTATTTACTCAGATTTTGTCTGATTGGACAAGGTGGTTTTGCTTTTT	625	0
TTTAGGTACCTTTTTGGTAGAATGCACTGAGACAGCGTCAGTTCTTCAGA	630	10
ATGCAACTCAGGATTCACTAGTAATCCTTGACGAACTGGGCAGAGGAACT	635	
AGTACTTTCGATGGATACGCCATTGCATACTCGGTAACCTGCTCTTCTCC	640	
TTCAACTTATACTTGTTGATCAACAAAAACATGCAATTCATTTTGCTGAA	645	
ACTTATTGATTTATATCAGGTTTTTCGTCACCTGGTAGAGAAGTTCAAT	650	
GTCGGATGCTCTTTGCAACACATTACCACCCTCTCACCAAGGAATTCGCG	655	
TCTCACCCACGTGTCACCTCGAAACACATGGCTTGCGCATTCAAATCAAG	660	
Figure 11 (Continued)	155	-

Figure 11 (Continued)

20 / 28

ATCTGATTATCAACCACGTGGTTGTGATCAAGACCTAGTGTTCTTGTACC	6650
GTTTAACCGAGGGAGCTTGTCCTGAGAGCTACGGACTTCAAGTGGCACTC	6700
ATGGCTGGAATACCAAACCAAGTGGTTGAAACAGCATCAGGTGCTGCTCA	6750
AGCCATGAAGAGATCAATTGGGGAAAACTTCAAGTCAAG	6800
CTGAGTTCTCAAGTCTGCATGAAGACTGGCTCAAGTCATTGGTGGGTATT	6850
TCTCGAGTCGCCCACAACAATGCCCCCATTGGCGAAGATGACTACGACAC	6900
TTTGTTTTGCTTATGGCATGAGATCAAATCCTCTTACTGTGTTCCCAAAT	6950
AAATGGCTATGACATAACACTATCTGAAGCTCGTTAAGTCTTTTGCTTCT	7000
CTGATGTTTATTCCTCTTAAAAAATGCTTATATATCAAAAAATTGTTTCC	7050
TCGATTATAACAAGATTATATGTATCTGTCGGTTTAGCTATGGTATAT	7100
AATATATGTATGTTCATGAGATTGGTCAAGAGAAATACTCACAAACAGTA	7150
TATTAAGAAGGAAATATGTTTATGCATTAATTTAAGTTTCAAGATAAACT	7200
GCAAATAACCTCGACTAAAGTTGCAAAGACCAAACACAAATTACAAAACT	7250
TATAAGACTTAAGTTCTGAATTCCCTAAAAACCAAAAAAAA	7300
TATTTTGTTGCATCTACAAACAACACAAACCTACATAGTTTATAACTTAC	7350
TCATCACTGAGATTAACATCAGAATCATTCTCCATTTCTTCATCTTCACT	7400
CTCATCATCACCACCACCATGATGATTCTCCTCCTCTTCACGTAACC	7450
TAGCAATCTCACTCTGAGCTCTATCAACAATCTGCTTCTTCTGCAACTCC	7500
AAATCTCTCTGAAAATCAGCTCTCATCTTCTCCAACTCCTTCATTTGCTC	7550
TTTCTTACTCTTCTCCATCTTCTCATAAACCTTCCCAAACCTCTCAACAG	7600
AATCCGCCAACATCTTATACGAAGCAGCGTCATTAACCTTCTTCCTCTCG	7650
TACTCAACCTCATCCTCATCCTCCTCCTCTTCAGAATCACCAGGACT	7700
ATCCATCATCTCATCAAACCCATTAGACTTATCTAAATAAA	7750
TCATAAACACAAACTCACCTGAATCAACACCACAAGCTAAACCTAAATCC	7800
GACTTGGGCGAAACACAAAGCAACATATCCAACTTATTGAAAAACGACCA	7850
TTTACTTGAACCTAAACCTGATTTCTCAACCTTAATCTTCTCTTTTCTAT	7900
ACTTCCTCTTCAAGTCATCAATCATTCTCCTACATTGCGTCTCAGATTTC	7950
TCCATCCTTAGCTCCTCACTCACTTCTCAGCTACTTCATTCCAATCCTC	8000
GTTCCTCAAACTCCTTCTACCCAATTGCAAAAACCTATCTCCCCAAACTT	8050
CAAGCAACACAA	8062

Figure 11 (Continued)




Figure 12

Comments/References: AtMSH3 3' side antisense : AtMSH3 3' (13 = 2104bp) from pUC18/13 Sal1/Sst1/T4 into pCW164 BarnH1/T4 in Agrobacterium LBA44O4




Figure 13

Comments/References: AtMSH6 (S8) 3' side antisens : 62 Sal1/Sst1/T4 (1379bp) into pCW164 BamH1/T4

422 Rec'd PST/PTO 1 0 APR 2000




Figure 14

Comments/References: AtMSH6 3'/AtMSH3 3' antisense : AtMSH6 (S8) 3' side (62=1379bp) Sal1/Sst1/T4 into pPF13 (pCW164 AtMSH3 (S5) 3' side (13=2104) antisens)/Sma1. in LBA4404

,1

••

WO 99/19492




Figure 15

Comments/References: AtMSH3 3'/AtMSH6 3' antisens (D): AtMSH3 (S5) 3' side (13=2104bp) Sal1/Sst1/T4 into pPF14 (AtMSH6 (S8) 3'side (62=1379bp) antisense into pCW164)/Sma1. in LBA4404




Figure 16

Comments/References: AtMSH3 (S8) complete, sense orientation : pPF26 (3342bp)

Sma1 into pCW164 Sma1

422 Rec'd PCT/PTO 1 0 APR 2000




Figure 17

Comments/References: pPZP111 with codA EcoR1 cassette in EcoR1 site and Hind3 GUS cassette in Hind3 efte. KenR. All genes under Promoter/terminator 35S

FR ALL OF BEET

Figure 18





Figure 19

p3243



SEQUENCE LISTING

<110>	Rhone-Poulenc Agro; Betzner, Andreas Stefan; Doutriaux, Marie-Pascale; Freyssinet, Georges; Perez, Pascual.	
<120>	Methods for obtaining plant varieties	
	in the second se	
<130>	395498C	
<150>	PO9745	`
<151>	1997-10-10	
<160>	98	
<210>	1	
<211>	23	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<221>	modified_base	
<222>	11	
<223>	I	
<220>		
<221>	modified_base	
<222>	14	
<223>	I	
<220>		
<221>	modified_base	
<222>	17	
<223>	I	
(223)	•	
<220>		
<223>	Degenerate oligonucleotides UPMU used to isolate AtMSH3 a AtMSH6.	ind
<300>		
<301>	Reenan and Kolodner	
<302>	Genetics	
<303>	132	
<306>	963-973	
<307>	1992	
<400>	1	

ctggatccac nggnccnaay atg

<210> 2 <211> 23 <212> DNA



```
<213>
             Artificial sequence
<220>
            modified base
<221>
<222>
             15
<223>
             Ι
<220>
<221>
             modified base
<222>
             18
<223>
             I
<220>
<223>
             Degenerate oligonucleotides DOMU used to isolate AtMSH3 and
             AtMSH6.
<300>
<301>
             Reenan and Kolodner
<302>
             Genetics
<303>
             132
<306>
             963-973
<307>
             1992
<400>
ctggatccrt artgngtnrc raa
                                                                       23
<210>
             3
<211>
             24
<212>
             DNA
<213>
             Artificial sequence
<220>
<223>
             MSH3 specific primer 636 for PCR using cDNA of Arabidopsis
             thaliana ecotype Columbia
<400>
tgctagtgcc tcttgcaagc tcat
                                                                        24
<210>
             4
<211>
             27
<212>
             DNA
<213>
             Artificial sequence
<220>
<223>
             Primer AP1 for PCR using cDNA of Arabidopsis thaliana ecotype
             Columbia containing adapter sequences ligated to both its
             ends
<400>
              4
```

						2	
		, / <u>1</u>					•
	•						
					**		
						4,	
				4.0			
a):							
			*				

	3	
ccatcctaat a	cgactcact atagggc	27
<210>	. 5	
<211>	23	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Primer AP2 for PCR using cDNA of Arabidopsis thaliana Columbia containing adapter sequences ligated to both ends	ecotype its
< 400>	5	
actcactata q	gggctcgagc ggc	23
<210>	6	
<211>	30	
<212>	DNA	
<213>	Artificial sequence	
<2137	Altificial sequence	
<220>		
<223>	MSH3 specific primer S525 for PCR using cDNA of Arabi thaliana ecotype Columbia	dopsis
<400>	6	
aggttctgat	tatgtgtgac gctttactta	30
<210>	7	
<211>	29	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	MSH3 specific primer S51 for PCR using cDNA of Arabic thaliana ecotype Columbia	lopsis
<400>	7 -	
ggatcgggta	ctgggttttg agtgtgagg	29
<210>	8	
<211>	24	
<212>	DNA	
<213>	Artificial sequence	
	Wittigar seducine	
<220>		
<223>	MSH3 specific primer 635 for PCR using cDNA of Arabi thaliana ecotype Columbia	dopsis

			-

<400>	8	
gcacgtgctt	gatggtgttt tcac 24	4
<210 > <211 > <212 > <213 >	9 28 DNA Artificial sequence	
<220> <223>	MSH3 specific primer S523 for PCR using cDNA of Arabidops thaliana ecotype Columbia	is
<400>	9	
tcagacagta	tccagcatgg cagaagta 2	8
<210 > <211 > <212 > <212 > <213 >	10 33 DNA Artificial sequence	
<220> <223>	MSH3 specific primer 1S5 for PCR using cDNA of Arabidopsi thaliana ecotype Columbia	s
<400>	10	
atcccgggat	gggcaagcaa aagcagcaga cga 3	3
<210> <211> <212> <213>	11 27 DNA Artificial sequence	
<220> <223>	MSH3 specific primer S53 for PCR using cDNA of Arabidopsi thaliana ecotype Columbia	.s
<400>	11	
gacaaagagc	gaaatgaggc cccttgg 2	7
<210> <211> <212> <213>	12 1250 DNA Arabidopsis thaliana ecotype Columbia	
<223>	Clone 52	

		4	

WO 99/19492 PCT/EP98/06977

cccgggatgg gcaago	caaaa gcagcagacg	atttctcgtt	tcttcgctcc	caaacccaaa	60
tccccgactc acgaac	ccgaa tccggtagcc	gaatcatcaa	caccgccacc	gaagatatcc	120
gccactgtat ccttct	tctcc ttccaagcgt	aagcttctct	ccgaccacct	cgccgccgcg	180
tcacccaaaa agccta	aaact ttctcctcac	actcaaaacc	cagtacccga	tcccaattta	240
caccaaagat ttctcc	cagag atttctggaa	ccctcgccgg	aggaatatgt	tcccgaaacg	300
tcatcatcga ggaaat	tacac accattggaa	cagcaagtgg	tggagctaaa	gagcaagtac	360
ccagatgtgg ttttga	atggt ggaagttggt	tacaggtaca	gattcttcgg	agaagacgcg	420
gagatogoag caegog	gtgtt gggtatttac	gctcatatgg	atcacaattt	catgacggcg	480
agtgtgccaa cattto	cgatt gaatttccat	gtgagaagac	tggtgaatgc	aggatacaag	540
attggtgtag tgaago	cagac tgaaactgca	gccattaagt	cccatggtgc	aaaccggacc	600
ggcccttttt tccggg	ggact groggogits	tataccaaag	ccacgcttga	agcggctgag	660
gatataagtg gtggtt	tgtgg tggtgaagaa	ggttttggtt	cacagagtaa	tttcttggtt	720
tgtgttgtgg atgaga	agagt taagtoggag	g acattagget	gtggtattga	aatgagtttt	780
gatgttagag teggte	gttgt tggcgttgaa	atttcgacag	gtgaagttgt	ttatgaagag	840
ttcaatgata atttca	atgag aagtggatta	a gaggetgtga	ttttgagctt	gtcaccagct	900
gagetgttge ttgged	cagee tettteacaa	caaactgaga	agtttttggt	ggcacatgct	960
ggacctacct caaacq	gttcg agtggaacgt	gcctcactgg	attgtttcag	caatggtaat	1020
gcagtagatg aggtta	atttc attatgtgaa	a aaaatcagcg	caggtaactt	agaagatgat	1080
aaagaaatga agctg	gaggc tgctgaaaaa	a ggaatgtett	gcttgacagt	tcatacaatt	1140
atgaacatgc cacato	ctgac tgttcaagco	ctcgccctaa	cgttttgcca	tctcaaacag	1200
tttggatttg aaagga	atect ttaccaagge	g gcctcatttc	gctctttgtc		1250

<210> 13
<211> 34
<212> DNA
<213> Artificial sequence

<400>

12

<220>
<223> MSH3 specific primer 2S5 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia

		٤,		7 ()	
					٠
			ii.		
90					
			*		
		•			

WO 99/19492 PCŢ/EP98/06977

(400>	13	•				
atcoogggto a	aaatgaaca a	agttggtttt	agtc			34
<210>	14					
<211>	27					
<212>	DNA					
<213>	Artificial	L sequence				
<220>						
<223>	-	ific primer ecotype Col		R using cDN	A of Arabido	psis
< 400>	14					
gccacatctg	actgttcaag	ccctcgc				27
22.0						
<210>	15					
<211>	2110					
<212>	DNA					
<213>	Arabidops	is thaliana	ecotype Co	lumbia		
<223>	Clone 13					
< 400>	15					
gccacatctg	actgttcaag	ccctcgccct	aacgttttgc	catctcaaac	agtttggatt	60
tgaaaggatc	ctttaccaag	gggcctcatt	tegetetttg	tcaagtaaca	cagagatgac	120
tctctcagcc	aatactctgc	aacagttgga	ggttgtgaaa	aataattcag	atggatcgga	180
atctggctcc	ttattccata	atatgaatca	cacacttaca	gtatatggtt	ccaggettet	240
tagacactgg	gtgactcatc	ctctatgcga	tagaaatttg	atatetgete	ggcttgatgc	300
tgtttctgag	atttctgctt	gcatgggatc	tcatagttct	tcccagetca	gcagtgagtt	360
ggttgaagaa	ggttctgaga	gagcaattgt	atcacctgag	ttttatctcg	tgctctcctc	420
agtcttgaca	gctatgtcta	gatcatctga	tattcaacgt	ggaataacaa	gaatctttca	480
tcggactgct	aaagccacag	agttcattgc	agttatggaa	gctattttac	ttgcggggaa	540
gcaaattcag	cggcttggca	taaagcaaga	ctctgaaatg	aggagtatgc	aatctgcaac	600
tgtgcgatct	actcttttga	gaaaattgat	ttctgttatt	tcatcccctg	ttgtggttga	660
caatgccgga	aaacttctct	ctgccctaaa	taaggaagcg	gctgttcgag	gtgacttgct	720
cgacatacta	atcacttcca	gcgaccaatt	teetgagett	gctgaagctc	gccaagcagt	780
tttagtcatc	agggaaaagc	tggattcctc	gatagcttca	tttcgcaaga	agctcgctat	840

			- 1/2	
	Ť			į

7.

tcgaaatttg	gaatttette	aagtgtcggg	gatcacacat	ttgatagagc	tgcccgttga	900
ttccaaggtc	cctatgaatt	gggtgaaagt	aaatagcacc	aagaagacta	ttcgatatca	960
tcccccagaa	atagtagctg	gcttggatga	gctagctcta	gcaactgaac	atcttgccat	1020
tgtgaaccga	gcttcgtggg	atagtttcct	caagagtttc	agtagatact	acacagattt	1080
taaggctgcc	gttcaagctc	ttgctgcact	ggactgtttg	cactcccttt	caactctatc	1140
tagaaacaag	aactatgtcc	gtcccgagtt	tgtggatgac	tgtgaaccag	ttgagataaa	1200
catacagtct	ggtcgtcatc	ctgtactgga	gactatatta	caagataact	tcgtcccaaa	1260
tgacacaatt	ttgcatgcag	aaggggaata	ttgccaaatt	atcaccggac	ctaacatggg	1320
aggaaagagc	tgctatatcc	gtcaagttgc	tttaatttcc	ataatggctc	aggttggttc	1380
ctttgtacca	gcgtcattcg	ccaagetgca	cgtgcttgat	ggtgttttca	ctcggatggg	1440
tgcttcagac	agtatccagc	atggcagaag	tacctttcta	gaagaattaa	gtgaagcgtc	1500
acacataatc	agaacctgtt	cttctcgttc	gcttgttata	ttagatgagc	ttggaagagg	1560
cactagcaca	cacgacggtg	tagccattgc	ctatgcaaca	ttacagcatc	tcctagcaga	1620
aaagagatgt	ttggttcttt	ttgtcacgca	ttaccctgaa	atagctgaga	tcagtaacgg	1680
attcccaggt	tctgttggga	cataccatgt	ctcgtatctg	acattgcaga	aggataaagg	1740
cagttatgat	catgatgatg	tgacctacct	atataagctt	gtgcgtggtc	tttgcagcag	1800
gagctttggt	tttaaggttg	ctcagcttgc	ccagatacct	ccatcatgta	tacgtcgagc	1860
catttcaatg	gctgcaaaat	tggaagctga	ggtacgtgca	agagagagaa	atacacgcat	1920
gggagaacca	gaaggacatg	aagaaccgag	aggcgcagaa	gaatctattt	cggctctagg	1980
tgacttgttt	gcagacctga	aatttgctct	ctctgaagag	gacccttgga	aagcattcga	2040
gtttttaaag	catgcttgga	agattgctgg	caaaatcaga	ctaaaaccaa	cttgttcatt	2100
ttgacccggg						2110

<210>	16
<211>	29
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	MSH3 specific primer S51 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia

<400>

			7	
4				

ggatcgggta	tgggttttg a g tgtgagg		. 29
<210> <211> <212> <213>	17 30 DNA Artificial sequence		
<220>		,	une.
<223>	MSH3 specific primer S52 thaliana ecotype Columbi	-	f Arabidopsis
<400>	17		
aggttctgat	tatgtgtgac gctttactta		30
<210><211><211><212><213>	18 3522 DNA Arabidopsis thaliana eco	otype Columbia	
<220><221><222><223>	CDS (100)(3342) AtMSH3 full-length cDNA polypeptide	and deduced sequence o	of the encoded
<400>	18		
cctaagaaag	cgcgcgaaaa ttggcaaccc aagt	tegeca tagecaegae caeç	gacette 60
catttctctt	aaacggagga gattacgaat aaag	JCaatt	9 9
	caa aag cag cag acg att t Gln Lys Gln Gln Thr Ile S 5	_	Lys
	ccg act cac gaa ccg aat o Pro Thr His Glu Pro Asn I 20 25		
	aag ata tcc gcc act gta t Lys Ile Ser Ala Thr Val 8 40		
	tee gae cae ete gee gee g . Ser Asp His Leu Ala Ala 2 55		
	cac act caa aac cca gta His Thr Gln Asn Pro Val	-	

	÷	,			
,			4		
					•

									•								
							gaa Glu									31	87
							tac Tyr									4:	35
							gat Asp 120									4:	83
							gaa Glu									5	31
							gat Asp									5	79
							cat His									6	27
							cag Gln									6	75
							cct Pro 200					_	_			7	23
							gcg Ala									7	71
							tca Ser									8	19
							gag Glu									8	67
agt Ser	ttt Phe	gat Asp	gtt Val 260	aga Arg	gtc Val	ggt Gly	gtt Val	gtt Val 265	ggc	gtt Val	gaa Glu	att Ile	tcg Ser 270	aca Thr	ggt Gly	9	15
							aat Asn 280									9	63

a.	-		
			•

	•				-	_			_		_	-		ggc		1011
											_			gga Gly		1059
														agc Ser 335		1107
			_	-		-				-	-			agc Ser	_	1155
			_	_	_		_	-	_	_		_	-	gaa Glu		1203
			_	_						-		_		cat His	_	1251
	_		-		-		-		_				_	ttt Phe		1299
								_			_		-	tca Ser 415	_	1347
			_				-			_		_	_	gag Glu	~	1395
gtg Val	aaa Lys	aat Asn 435	aat Asn	tca Ser	gat Asp	gga Gly	tcg Ser 440	gaa Glu	tct Ser	ggc Gly	tcc Ser	tta Leu 445	ttc Phe	cat His	aat Asn	1443
														cac His		1491
												_		ctt Leu		1539
					Ser					Ser				tcc Ser 495		1587

. ,	•		
			-
			•

									• .							
ccc	agc Ser	agt Ser	gag Glu 500	ttg Leu	grt Val	gaa Glu	gaa Glu	ggt Gly 505	tct Ser	gag Glu	aga Arg	gca Ala	att Ile 510	gta Val	tca Ser	1635
cct Pro	gag Glu	ttt Phe 515	tat Tyr	ctc Leu	gtg Val	ctc Leu	tcc Ser 520	tca Ser	gtc Val	ttg Leu	aca Thr	Ala 525	atg Met	tct Ser	aga Arg	1683
tca Ser	tct Ser 530	gat Asp	att Ile	caa Gln	cgt Arg	gga Gly 535	ata Ile	aca Thr	aga Arg	atc Ile	ttt Phe 540	cat His	cgg Arg	act Thr	gct Ala	1731
aaa Lys 545	gcc Ala	aca Thr	gag Glu	ttc Phe	att Ile 550	gca Ala	gtt Val	atg Met	gaa Glu	gct Ala 555	att Ile	tta Leu	ctt Leu	gcg Ala	260 GJA aaa	1779
aag Lys	caa Gln	att Ile	cag Gln	cgg Arg 565	ctt Leu	ggc Gly	ata Ile	aag Lys	caa Gln 570	gac Asp	tct Ser	gaa Glu	atg Met	agg Arg 575	agt Ser	1827
atg Met	caa Gln	tct Ser	gca Ala 580	act Thr	gtg Val	cga Arg	tct Ser	act Thr 585	ctt Leu	ttg Leu	aga Arg	aaa Lys	ttg Leu 590	att Ile	tct Ser	1875
gtt Val	att Ile	tca Ser 595	tcc Ser	cct Pro	gtt Val	gtg Val	gtt Val 600	gac Asp	aat Asn	gcc Ala	gga Gly	aaa Lys 605	ctt Leu	ctc Leu	tct Ser	1923
gcc Ala	cta Leu 610	aat Asn	aag Lys	gaa Glu	gcg Ala	gct Ala 615	gtt Val	cga Arg	ggt Gly	gac Asp	ttg Leu 620	ctc Leu	gac Asp	ata Ile	cta Leu	1971
atc Ile 625	act Thr	tcc Ser	agc Ser	gac Asp	caa Gln 630	ttt Phe	cct Pro	gag Glu	ctt Leu	gct Ala 635	gaa Glu	gct Ala	cgc Arg	caa Gln	gca Ala 640	2019
gtt Val	tta Leu	gtc Val	atc Ile	agg Arg 645	gaa Glu	aag Lys	ctg Leu	gat Asp	tcc Ser 650	tcg Ser	ata Ile	gct Ala	tca Ser	ttt Phe 655	cgc Arg	2067
aag Lys	aag Lys	ctc Leu	gct Ala 660	att Ile	cga Arg	aat Asn	ttg Leu	gaa Glu 665	ttt Phe	ctt Leu	caa Gln	gtg Val	tcg Ser 670	GJA aaa	atc Ile	2115
aca Thr	cat His	ttg Leu 675	ata Ile	gag Glu	ctg Leu	ccc Pro	gtt Val 680	gat Asp	tcc Ser	aag Lys	gtc Val	cct Pro 685	atg His	aat Asn	tgg Trp	2163
gtg Val	aaa Lys 690	gta Val	aat Asn	agc Ser	acc Thr	aag Lys 695	aag Lys	act Thr	att Ile	cga Arg	tat Tyr 700	cat His	ccc Pro	cca Pro	g aa Glu	2211

	•		
			. 4;
			•

								gct Ala								22!	59
att Ile	gtg Val	aac Asn	cga Arg	gct Ala 725	tcg Ser	tgg Trp	gat Asp	agt Ser	ttc Phe 730	ctc Leu	aag Lys	agt Ser	ttc Phe	agt Ser 735	aga Arg	230	07
tac Tyr	tac Tyr	aca Thr	gat Asp 740	ttt Phe	aag Lys	gct Ala	gcc Ala	gtt Val 745	caa Gln	gct Ala	ctt Leu	gct Ala	gca Ala 750	ctg Leu	gac Asp	23	55
tgt Cys	ttg Leu	cac His 755	tcc Ser	ctt Leu	tca Ser	act Thr	cta Leu 760	tct Ser	aga Arg	aac Asn	aag Lys	aac Asn 765	tat Tyr	gtc Val	cgt Arg	240	03
ccc Pro	gag Glu 770	ttt Phe	gtg Val	gat Asp	gac Asp	tgt Cys 775	gaa Glu	cca Pro	gtt Val	gag Glu	ata Ile 780	aac Asn	ata Ile	cag Gln	tct Ser	24!	51
ggt Gly 785	cgt Arg	cat His	cct Pro	gta Val	ctg Leu 790	gag Glu	act	ata Ile	tta Leu	caa Gln 795	gat Asp	aac Asn	ttc Phe	gtc Val	cca Pro 800	245	99
aat Asn	gac Asp	aca Thr	att Ile	ttg Leu 805	cat His	gca Ala	gaa Glu	gly aaa	gaa Glu 810	tat Tyr	tgc Cys	caa Gln	att Ile	atc Ile 815	acc Thr	254	47
gga Gly	cct Pro	aac Asn	atg Met 820	gga Gly	gga Gly	aag Lys	agc Ser	tgc Cys 825	tat Tyr	atc Ile	cgt Arg	caa Gln	gtt Val 830	gct Ala	tta Leu	259	95
att Ile	tcc Ser	ata Ile 835	atg Met	gct Ala	cag Gln	gtt Val	ggt Gly 840	tcc Ser	ttt Phe	gta Val	cca Pro	gcg Ala 845	tca Ser	ttc Phe	gcc Ala	264	43
aag Lys	ctg Leu 850	cac His	gtg Val	ctt Leu	gat Asp	ggt Gly 855	gtt Val	ttc Phe	act Thr	cgg Arg	atg Met 860	ggt Gly	gct Ala	tca Ser	gac Asp	269	91
agt Ser 865	atc Ile	cag Gln	cat His	ggc Gly	aga Arg 870	agt Ser	acc Thr	ttt Phe	cta Leu	gaa Glu 875	gaa Glu	tta Leu	agt Ser	gaa Glu	gcg Ala 880	273	39
tca Ser	cac His	ata Ile	atc I le	aga Arg 885	acc Thr	tgt Cys	tct Ser	tct Ser	cgt Arg 890	tcg Ser	ctt Leu	gtt Val	ata Ile	tta Leu 895	gat Asp	278	87
gag Glu	ctt Leu	gga Gly	aga Arg 900	ggc Gly	act Thr	agc Ser	aca Thr	cac His 905	gac Asp	ggt Gly	gta Val	gcc Ala	att Ile 910	gcc Ala	tat Tyr	283	35

			•
		*	
	•		•
4			
		,,	
•			
1			
•	*.		

13	
gca aca tta cag cat ctc cta gca gaa aag aga tgt ttg gtt ctt ttt Ala Thr Leu Gln His Leu Leu Ala Glu Lys Arg Cys Leu Val Leu Phe 915 920 925	2883
gtc acg cat tac cct gaa ata gct gag atc agt aac gga ttc cca ggt Val Thr His Tyr Pro Glu Ile Ala Glu Ile Ser Asn Gly Phe Pro Gly 930 935 940	2931
tct gtt ggg aca tac cat gtc tcg tat ctg aca ttg cag aag gat aaa Ser Val Gly Thr Tyr His Val Ser Tyr Leu Thr Leu Gln Lys Asp Lys 945 950 955 960	2979
ggc agt tat gat cat gat gtg acc tac cta tat aag ctt gtg cgt Gly Ser Tyr Asp His Asp Asp Val Thr Tyr Leu Tyr Lys Leu Val Arg 965 970 975	3027
ggt ctt tgc agc agg agc ttt ggt ttt aag gtt gct cag ctt gcc cag Gly Leu Cys Ser Arg Ser Phe Gly Phe Lys Val Ala Gln Leu Ala Gln 980 985 990	3075
ata cct cca tca tgt ata cgt cga gcc att tca atg gct gca aaa ttg Ile Pro Pro Ser Cys Ile Arg Arg Ala Ile Ser Met Ala Ala Lys Leu 995 1000 1005	3123
gaa gct gag gta cgt gca aga gag aga aat aca cgc atg gga gaa cca Glu Ala Glu Val Arg Ala Arg Glu Arg Asn Thr Arg Met Gly Glu Pro 1010 1015 1020	3171
gaa gga cat gaa gaa ccg aga ggc gca gaa gaa tct att tcg gct cta Glu Gly His Glu Glu Pro Arg Gly Ala Glu Glu Ser Ile Ser Ala Leu 1025 1030 1035 1040	3219
ggt gac ttg ttt gca gac ctg aaa ttt gct ctc tct gaa gag gac cct Gly Asp Leu Phe Ala Asp Leu Lys Phe Ala Leu Ser Glu Glu Asp Pro 1045 1050 1055	3267
tgg aaa gca ttc gag ttt tta aag cat gct tgg aag att gct ggc aaa Trp Lys Ala Phe Glu Phe Leu Lys His Ala Trp Lys Ile Ala Gly Lys 1060 1065 1070	3315
atc aga cta aaa cca act tgt tca ttt tgatttaatc ttaacattat Ile Arg Leu Lys Pro Thr Cys Ser Phe 1075 1080	3362
agcaactgca aggtettgat catetgttag ttgegtaeta aettatgtgt attagtataa	3422
caagaaaaga gaattagaga gatggattet aateeggtgt tgeagtaeat etttteteea	3482
cccgcataaa aaaaaaaaa aaaaaaaaaaaaaaaaaaa	3522

<210> 19 <211> 1081 <212> PRT

		¢.	
÷			
			•
			•

Arabidopsis thaliana ecotype Columbia Polypeptide MSH3 <223> <400> 19 Met Gly Lys Gln Lys Gln Gln Thr Ile Ser Arg Phe Phe Ala Pro Lys Pro Lys Ser Pro Thr His Glu Pro Asn Pro Val Ala Glu Ser Ser Thr 20 25 Pro Pro Pro Lys Ile Ser Ala Thr Val Ser Phe Ser Pro Ser Lys Arg 40 35 Lys Leu Leu Ser Asp His Leu Ala Ala Ala Ser Pro Lys Lys Pro Lys Leu Ser Pro His Thr Gln Asn Pro Val Pro Asp Pro Asn Leu His Gln Arg Phe Leu Gln Arg Phe Leu Glu Pro Ser Pro Glu Glu Tyr Val Pro 90 85 Glu Thr Ser Ser Ser Arg Lys Tyr Thr Pro Leu Glu Gln Gln Val Val 100 105 Glu Leu Lys Ser Lys Tyr Pro Asp Val Val Leu Met Val Glu Val Gly Tyr Arg Tyr Arg Phe Phe Gly Glu Asp Ala Glu Ile Ala Ala Arg Val 135 140 Leu Gly Ile Tyr Ala His Met Asp His Asn Phe Met Thr Ala Ser Val 145 150 155 160 Pro Thr Phe Arg Leu Asn Phe His Val Arg Arg Leu Val Asn Ala Gly 165 170 Tyr Lys Ile Gly Val Val Lys Gln Thr Glu Thr Ala Ala Ile Lys Ser 185 His Gly Ala Asm-Arg Thr Gly Pro Phe Phe Arg Gly Leu Ser Ala Leu 195 200 205 Tyr Thr Lys Ala Thr Leu Glu Ala Ala Glu Asp Ile Ser Gly Gly Cys 210 215 220 Gly Glu Glu Gly Phe Gly Ser Gln Ser Asn Phe Leu Val Cys Val 225 230 235 Val Asp Glu Arg Val Lys Ser Glu Thr Leu Gly Cys Gly Ile Glu Met 250 245

			•

WO 99/19492

	Ser	Phe	Asp	Val 260	Arg	Val	Gly	Val	Val 265	Gly	Val	Glu	Ile	Ser 270	Thr	Gly
•	Glu	Val	Val 275	Tyr	Glu	Glu	Phe	Asn 280	Asp	Asn	Phe	Met	Arg 285	Ser	Gly	Leu
•	Glu	Ala 290	Val	Ile	Leu	Ser	Leu 295	Ser	Pro	Ala	Glu	Leu 300	Leu	Leu	Gly	Gln
	Pro 305	Leu	Ser	Gln	Gln	Thr 310	Glu	Lys	Phe	Leu	Val 315	Ala	Met	Ala	Gly	Pro 320
•	Thr	Ser	Asn	Val	Arg 325	Val	Glu	Arg	Ala	Ser 330	Leu	Asp	Cys	Phe	Ser 335	Asn
	Gly	Asn	Ala	Val 340	Asp	Glu	Val	Ile	Ser 345	Leu	Cys	Glu	Lys	Ile 350	Ser	Ala
	Gly	Asn	Leu 355	Glu	Asp	Asp	Lys	Glu 360	Met	Lys	Leu	Glu	Ala 365	Ala	Glu	Lys
	Gly	Мет 370	Ser	Cys	Leu	Thr	Val 375	His	Thr	Ile	Met	Asn 380	Met	Pro	His	Leu
	Thr 385	Val	Gln	Ala	Leu	Ala 390	Leu	Thr	Phe	Cys	His 395	Leu	Lys	Gln	Phe	Gly 400
	Phe	Glu	Arg	Ile	Leu 405	туr	Gln	Gly	Ala	Ser 410	Phe	Arg	Ser	Leu	Ser 415	Ser
	Asn	Thr	Glu	Met 420	Thr	Leu	Ser	Ala	Asn 425	Thr	Leu	Gln	Gln	Leu 430	Glu	Val
	Val	Lys	Asn 435	Asn	Ser	Asp	Gly	Ser 440	Glu	Ser	Gly	Ser	Leu 445	Phe	His	Asn
	Met	Asn 450	His	Thr	Leu	Thr	Val 455	Tyr	Gly	Ser	Arg	Leu 460	Leu	Arg	His	Trp
	Val 465	Thr	His	Pro	Leu	Cys 470	Asp	Arg	Asn	Leu	Ile 475	Ser	Ala	Arg	Leu	Asp 480
	Ala	Val	Ser	Glu	Ile 485	Ser	Ala	Cys	Met	Gly 490		His	Ser	Ser	Ser 495	Gln
	Leu	Ser	Ser	Glu 500	Leu	Val	Glu	Glu	Gly 505	Ser	Glu	Arg	Ala	Ile 510	Val	Ser
	Pro	Glu	Phe 515		Leu	Val	Leu	Ser 520	Ser	Val	Leu	Thr	Ala 525	Met	Ser	Arg
	Ser	Ser	Asp	Ile	Gln	Arg	Gly	Ile	Thr	Arg	Ile	Phe	His	Arg	Thr	Ala

			,1	
			ē.	
,				
		•		

Lys 545	Ala	Thr	Glu	Phe	11e 550	Ala	Val	Met	Glu	Ala 555	Ile	Leu	Leu	Ala	Gly 560
Lys	Gln	Ile	Gln	Arg 565	Leu	Gly	Ile	Lys	Gln 570	Asp	Ser	Glu	Met	Arg 575	Ser
Met	Gln	Ser	Ala 580	Thr	Val	Arg	Ser	Thr 585	Leu	Leu	Arg	Lys	Leu. 590	Ile	Ser
Val	Ile	Ser 595	Ser	Pro	Val	Val	Val 600	Asp	Asn	Ala	Gly	Lys 605	Leu	Leu	Ser
Ala	Leu 610	Asn	Lys	Glu	Ala	Ala 615	Val	Arg	Gly	Asp	Leu 620	Leu	Asp	Ile	Leu
Ile 625	Thr	Ser	Ser	Asp	Gln 630	Phe	Pro	Glu	Leu	Ala 635	Glu	Ala	Arg	Gln	Ala 640
Val	Leu	Val	Ile	Arg 645	Glu	Lys	Leu	Asp	Ser 650	Ser	Ile	Ala	Ser	Phe 655	Arg
Lys	Lys	Leu	Ala 660	Ile	Arg	Asn	Leu	Glu 665	Phe	Leu	Gln	Val	Ser 670	Gly	Ile
		675	Ile				680	-		_		685	-		-
	690		Asn			695	_				700				
705			Gly		710					715					720
			Arg	725		-	-		730					735	
			Asp 740					745					750		
Cys	Leu	His 755	Ser	Leu	Ser	Thr	Leu 760	Ser	Arg	Asn	Lys	Asn 765	Tyr	Val	Arg
Pro	Glu 770	Phe	Val	Asp	Asp	Cys 775	Glu	Pro	Val	Glu	Ile 780	Asn	Ile	Gln	Ser
Gly 785	Arg	His	Pro	Val	Leu 790	Glu	Thr	Ile	Leu	Gln 795	Asp	Asn	Phe	Val	Pro 800
				805				-	810					815	Thr
Gly	Pro	Asn	Met 820	Gly	Gly	Lys	Ser	Cys 825		Ile	Arg	Gln	Val 830	Ala	Leu

		•	
*			

Ile Ser Ile Met Ala Gln Val Gly Ser Phe Val Pro Ala Ser Phe Ala
835
845

Lys Leu His Val Leu Asp Gly Val Phe Thr Arg Met Gly Ala Ser Asp 850 855 860

Ser Ile Gln His Gly Arg Ser Thr Phe Leu Glu Glu Leu Ser Glu Ala 865 870 875 880

Ser His Ile Ile Arg Thr Cys Ser Ser Arg Ser Leu Val Ile Leu Asp 885 890 895

Glu Leu Gly Arg Gly Thr Ser Thr His Asp Gly Val Ala Ile Ala Tyr 900 905 910

Ala Thr Leu Gln His Leu Leu Ala Glu Lys Arg Cys Leu Val Leu Phe 915 920 925

Val Thr His Tyr Pro Glu Ile Ala Glu Ile Ser Asn Gly Phe Pro Gly 930 935 940

Ser Val Gly Thr Tyr His Val Ser Tyr Leu Thr Leu Gln Lys Asp Lys 945 950 955 960

Gly Ser Tyr Asp His Asp Asp Val Thr Tyr Leu Tyr Lys Leu Val Arg 965 970 975

Gly Leu Cys Ser Arg Ser Phe Gly Phe Lys Val Ala Gln Leu Ala Gln 980 985 990

Ile Pro Pro Ser Cys Ile Arg Arg Ala Ile Ser Met Ala Ala Lys Leu 995 1000 1005

Glu Ala Glu Val Arg Ala Arg Glu Arg Asn Thr Arg Met Gly Glu Pro 1010 1015 1020

Glu Gly His Glu Glu Pro Arg Gly Ala Glu Glu Ser Ile Ser Ala Leu 1025 1030 1035 1040

Gly Asp Leu Phe Ala Asp Leu Lys Phe Ala Leu Ser Glu Glu Asp Pro 1045 1050 1055

Trp Lys Ala Phe Glu Phe Leu Lys His Ala Trp Lys Ile Ala Gly Lys 1060 1065 1070

Ile Arg Leu Lys Pro Thr Cys Ser Phe 1075 1080

<210> 20

<211> 24

<212> DNA

<213> Artificial sequence

	, (E)		
			· .
	÷		•

18

<220>	•	
<223>	MSH6 specific primer 638 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia	
<400>	20	
tctctaccag g	grgacgaaaa accg 2	24
	'	
<210>	21	
<211>	28	
<212> <213>	DNA Artificial sequence	
<713>	Artificial sequence	
<220>		
<223>	Primer S81 for PCR using cDNA of Arabidopsis thaliana ecoty Columbia	φe
<400>	21	
cgtcgcctt :	agcatecect teetteac	28
<210>	22	
<211>	30	
<212>	DNA	
<213>	Artificial sequence	
.220		
<220> <223>	MSH6 specific primer S823 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia	3
<400>	22	
gcttggcgca	tctaatagaa tcatgacagg	30
<210>	23	
<211>	24	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	MSH6 specific primer 637 for PCR using cDNA of Arabidopsis	
	thaliana ecotype Columbia	
<400>	23	
gacagcgtca	gttcttcaga atgc	24
<210>	24	
<211>	33	

DNA

<212>

		,	,		
,					
	¥.				
				1-2	

<213>	Artificial sequence	
<220>		
<223>	MSH6 specific primer 1SB for PCR using cDNA of Arabidopsisthaliana ecotype Columbia	S
<400>	24	
atcccgggat	gcagcgccag agatcgattt tgt	33
<210>	25	
<211>	27	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	MSH6 specific primer S83 for PCR using cDNA of Arabidopsi thaliana ecotype Columbia	s
<400>	25	
egetatetat	ggctgcttcg aattgag	27
<210>	26	
<211>	1385	
<212>	DNA	
<213>	Arabidopsis thaliana ecotype Columbia	
<223>	Clone 43	
<400>	26	
cccgggatgc	agegecagag ategatiting tetiteties aaaaaceeae ggeggegaet	60
acgaagggtt	tggtttccgg cgatgctgct agcggcgggg gcggcagcgg aggaccacga	120
tttaatgtga	aggaagggga tgctaaaggc gacgcttctg tacgttttgc tgtttcgaaa	180
tctgtcgatg	aggttagagg aacggatact ccaccggaga aggttccgcg tcgtgtcctg	240
ccgtctggat	traagccggc rgaarccgcc ggrgargerr cgrcccrgrr crccaatarr	300
atgcataagt	ttgtaaaagt cgatgatcga gattgttctg gagagaggag ccgagaagat	360
gttgttccgc	tgaatgattc atctctatgt atgaaggcta atgatgttat tcctcaattt	420
cgttccaata	atggtaaaac tcaagaaaga aaccatgctt ttagtttcag tgggagagct	480
gaacttagat	cagtagaaga tataggagta gatggcgatg ttcctggtcc agaaacacca	540
gggatgcgtc	cacgtgcttc tcgcttgaag cgagttctgg aggatgaaat gacttttaag	600
gaggataagg	ttcctgtatt ggactctaac aaaaggctga aaatgctcca ggatccggtt	660

	40			
				· ·
				•
		÷.		

tgtggagaga	agaaagaagt	aaacgaagga	accaaatttg	aatggcttga	gtcttctcga	720
atcagggatg	ccaatagaag	acgtcctgat	gatccccttt	acgatagaaa	gaccttacac	780
ataccacctg	atgttttcaa	gaaaatgtct	gcatcacaaa	agcaatattg	gagtgttaag	840
agtgaatata	tggacattgt	gcttttcttt	aaagtgggga	aattttatga	gctgtatgag	900
ctagatgcgg	aattaggtca	caaggagctt	gactggaaga	tgaccatgag	tggtgtggga	960
aaatgcagac	aggttggtat	ctctgaaagt	gggatagatg	aggcagtgca	aaagctatta	1020
gctcgtggat	ataaagttgg	acgaatcgag	cagctagaaa	catctgacca	agcaaaagcc	1080
agaggtgcta	atactataat	tccaaggaag	ctagttcagg	tattaactcc	atcaacagca	1140
agcgagggaa	acatcgggcc	tgatgccgtc	catcttcttg	ctataaaaga	gatcaaaatg	1200
gagctacaaa	agtgttcaac	tgtgtatgga	tttgcttttg	ttgactgtgc	tgccttgagg	1250
ttttgggttg	ggtccatcag	cgatgatgca	tcatgtgctg	ctcttggagc	gttattgatg	1320
caggittete	caaaggaagt	gttatatgac	agtaaagggc	tatcaagaga	agcacaaaag	1330
gctctaagga	aatatacgtt	gacagggtct	acggcggtac	agttggctcc	agtaccacaa	1440
gtaatggggg	atacagatgc	tgctggagtt	agaaatataa	tagaatctaa	cggatacttt	1500
aaaggttctt	ctgaatcatg	gaactgtgct	gttgatggtc	taaatgaatg	tgatgttgcc	1560
cttagtgctc	ttggagagct	aattaatcat	ctgtctaggc	taaagctaga	agatgtactt	1620
aagcatgggg	atatttttcc	ataccaagtt	tacaggggtt	gtctcagaat	tgatggccag	1680
acgatggtaa	atcttgagat	atttaacaat	agctgtgatg	gtggtccttc	agggaccttg	1740
tacaaatatc	ttgataactg	tgttagtcca	actggtaagc	gactcttaag	gaattggatc	1800
tgccatccac	tcaaagatgt	agaaagcatc	aataaacggc	ttgatgtagt	tgaagaattc	1860
acggcaaact	cagaaagtat	gcaaatcact	ggccagtatc	tecacaaact	tccagactta	1920
gaaagactgc	tcggacgcat	caagtctagc	gttcgatcat	cagectetgt	gttgcctgct	1980
cttctgggga	aaaaagtgct	gaaacaacga	gttaaagcat	ttgggcaaat	tgtgaaaggg	2040
ttcagaagtg	gaattgatct	gttgttggct	ctacagaagg	aatcaaatat	gatgagtttg	2100
ctttataaac	tctgtaaact	tcctatatta	gtaggaaaaa	gcgggctaga	gttatttctt	2160
tctcaattcg	aagcagccat	agatagcg				2188

<210> <211> <212> <213> <223>	27 1385 DNA Arabidops Clone 62	sis thaliana	a ecotype Co	olumbia		
<400>	27					
catcagcctc	tgtgttgcct	gctcttctgg	ggaaaaaagt	gctgaaacaa	cgagttaaag	60
catttgggca	aattgtgaaa	gggttcagaa	gtggaattga	tctgttgttg	gctctacaga	120
aggaatcaaa	tatgatgagt	ttgctttata	aactctgtaa	acttcctata	ttagtaggaa	180
aaagcgggct	agagttattt	ctttctcaat	tcgaagcagc	catagatagc	gactttccaa	240
attatcagaa	ccaagatgtg	acagatgaaa	acgctgaaac	tctcacaata	cttatcgaac	300
tttttatcga	aagagcaact	caatggtctg	aggtcattca	caccataagc	tgcctagatg	360
tcctgagatc	ttttgcaatc	gcagcaagtc	tetetgetgg	aagcatggcc	aggcctgtta	420
tttttcccga	atcagaagct	acagatcaga	atcagasaac	aaaagggcca	atacttaaaa	480
tccaaggact	atggcatcca	tttgcagttg	cagccgatgg	tcaattgcct	gttccgaatg	540
atatactcct	tggcgaggct	agaagaagca	gtggcagcat	tcatcctcgg	tcattgttac	600
tgacgggacc	aaacatgggc	ggaaaatcaa	ctcttcttcg	tgcaacatgt	ctggccgtta	660
tctttgccca	acttggctgc	tacgtgccgt	gtgagtcttg	cgaaatctcc	ctcgtggata	720
ctatcttcac	aaggettgge	gcatctgata	gaatcatgac	aggagagagt	acctttttgg	780
tagaatgcac	tgagacagcg	tcagttcttc	agaatgcaac	tcaggattca	ctagtaatcc	840
ttgacgaact	gggcagagga	actagtactt	tcgatggata	cgccattgca	tactcggttt	900
ttcgtcacct	ggtagagaaa	gttcaatgtc	ggatgetett	tgcaacacat	taccaccctc	960
tcaccaagga	attegegtet	cacccacgtg	tcacctcgaa	acacatggct	tgcgcattca	1020
aatcaagatc	tgattatcaa	ccacgtggtt	gtgatcaaga	cctagtgttc	ttgtaccgtt	1080
taaccgaggg	agcttgtcct	gagagctacg	gacttcaagt	ggcactcatg	gctggaatac	1140
caaaccaagt	ggttgaaaca	gcatcaggtg	ctgctcaagc	catgaagaga	tcaattgggg	1200
aaaacttcaa	gtcaagtgag	ctaagatctg	agttctcaag	tctgcatgaa	gactggctca	1260
agtcattggt	gggtatttct	cgagtcgccc	acaacaatgc	ccccattggc	gaagatgact	1320
acgacacttt	gttttgctta	tggcatgaga	tcaaatcctc	ttactgtgtt	cccaaataac	1380

•	Ž,			
	•	<u>.</u>		
	÷			
				,
	-			•
	9.			
		÷		

WO 99/19492	PCŢ/EP98/06977
WO 99/19492	PCŢ/EP98/0

ccggg	1385
.210.	
<210>	28
<211>	34
<212>	DNA
<213>	Artificial sequence
<220>	-
<223>	MSH6 specific primer 2S8 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia
< 400>	26
atcccgggtt	atttgggaac acagtaagag gatt 34
<210>	29
<211>	27
<212>	DNA
<213>	
(213)	Artificial sequence
<220>	
<223>	MSH6 specific primer S82 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia
<400>	29
gcgttcgatc	atcagcctct gtgttgc 27
<210>	3.0
<211>	3606
<212>	DNA
<213>	
(213)	Arabidopsis thaliana ecotype Columbia
<220>	
<221>	CDS
<222>	(142)(3468)
<223>	AtMSH6 full-length cDNA and deduced sequence of the encoded polypeptide
<400>	30
aaaagttgag	ccctgaggag tatcgtttcc gccatttcta cgacgcaagg cgaaaatttt 60
tggcgccaat	ctttcccccc tttcgaattc tctcagctca aaacatcgtt tctctctcac 120
tctctctcac	aattccaaaa a atg cag cgc cag aga tcg att ttg tct ttc 171 Met Gln Arg Gln Arg Ser Ile Leu Ser Phe 1 5 10

					•	
		-	•			•
	•					-
				'n		•
				25		
			*			

											_	-		ggc Gly 25		219
														gtg Val		267
			_				_		_	-		_	_	tcg Ser		315
														gtt Val		363
			_					_	_	_	_		_	gst Gly	_	411
														gtc Val 105		459
														ccg Pro		507
Asn	Asp	Ser 125	Ser	Leu	Cys	Met	Lys 130	Ala	Asn	Asp	Val	Ile 135	Pro	caa Gln	Phe	555
														agt Ser		603
	Gly							-	-	_			_	gat Asp		651
													-	tct Ser 185		699
				Leu					Thr		_		_	aag Lys	_	747
			Asp					Leu					Asp	ccg Pro		795

		, ·	
			·
			•

			aag Lys													843
			cga Arg													891
			aga Arg													939
			tca Ser 270													987
			ctt Leu													1035
			gaa Glu													1083
			gga Gly													1131
			gtg Val													1179
			cta Leu 3 5 0													1227
			cca Pro													1275
agc Ser	gag Glu 380	gga Gly	aac Asn	atc Ile	Gly ggg	cct Pro 385	gat Asp	gcc Ala	gtc Val	cat His	ctt Leu 390	ctt Leu	gct Ala	ata Ile	aaa Lys	1323
gag Glu 395	atc Ile	aaa Lys	atg Met	gag Glu	cta Leu 400	caa Gln	aag Lys	tgt Cys	tca Ser	act Thr 405	gtg Val	tat Tyr	gga Gly	ttt Phe	gct Ala 410	1371
ttt Phe	gtt Val	gac Asp	tgt Cys	gct Ala 415	gcc Ala	ttg Leu	agg Arg	ttt Phe	tgg Trp 420	gtt Val	Gly aaa	tcc Ser	atc Ile	agc Ser 425	gat Asp	1419

			. ·	
	-			

gat Asp	gca Ala	tca Ser	tgt Cys 430	gct Ala	gct Ala	ctt Leu	gga Gly	gcg Ala 435	tta Leu	ttg Leu	atg Met	cag Gln	gtt Val 440	tct Ser	cca Pro	1467
aag Lys	gaa Glu	gtg Val 445	tta Leu	tat Tyr	gac Asp	agt Ser	aaa Lys 450	ggg Gly	cta Leu	tca Ser	aga Arg	gaa Glu 455	gca Ala	caa Gln	aag Lys	1515
													cag Gln			1563
													gtt Val			1611
ata Ile	ata Ile	gaa Glu	tct Ser	aac Asn 495	gga Gly	tac Tyr	ttt Phe	aaa Lys	ggt Gly 500	tct Ser	tct Ser	gaa Glu	tca Ser	tgg Trp 505	aac Asn	1659
													agt Ser 520			1707
													gat Asp			1755
													tgt Cys			1803
													aat Asn			1851
													aac Asn			1899
													cat His 600			1947
aaa Lys	gat Asp	gta Val 605	gaa Glu	agc Ser	atc Ile	aat Asn	aaa Lys 610	cgg Arg	ctt Leu	gat Asp	gta Val	gtt Val 615	gaa Glu	gaa Glu	ttc Phe	1995
acg Thr	gca Ala 620	aac Asn	tca Ser	gaa Glu	agt Ser	atg Met 625	caa Gln	atc Ile	act	ggc	cag Gln 630	tat Tyr	ctc Leu	cac His	aaa Lys	2043

		,		
				•
			,	•
4				
\$v				

		-		_	agá Arg 640	•			_		_		_	_	_	2091
		-			ttg Leu				_							2139
	-	_		_	ttt Phe								_	_		2187
	_	_	-	_	gct Ala		_	_				-	_	_	_	2235
				_	aaa Lys					_			_			2283
					caa Gln 720				_		_	_	_			2331
		-			gat Asp	-		-	-							2379
			_		ttt Phe		-	-						_		2427
				_	tgc Cys		-	-	_	_			Āla		_	2475
_	-			-	gga Gly		_	-			_					2523
					cag Gln 800	Asn					Gly					2571
					cat His					Ala					Leu	2619
				Asp	ata Ile				Glu					Ser	ggc	2667

÷	-		
			•

agc Ser	att Ile	cat His 845	cct Pro	cgg Arg	tca Ser	ttg Leu	tta Leu 950	ctg Leu	acg Thr	gga Gly	cca Pro	aac Asn 855	atg Met	ggc Gly	gga Gly		2715
aaa Lys	tca Ser 860	act Thr	ctt Leu	ctt Leu	cgt Arg	gca Ala 865	aca Thr	tgt Cys	ctg Leu	gcc Ala	gtt Val 870	atc Ile	ttt Phe	gcc Ala	caa Gln		2763
ctt Leu 875	ggc Gly	tgc Cys	tac Tyr	gtg Val	ccg Pro 880	tgt Cys	gag Glu	tct Ser	tgc Cys	gaa Glu 885	atc Ile	tcc Ser	ct <i>c</i> Leu	gtg Val	gat Asp 890		2811
act Thr	atc Ile	ttc Phe	aca Thr	agg Arg 895	ctt Leu	ggc Gly	gca Ala	tct Ser	gat Asp 900	aga Arg	atc Ile	atg Met	aca Thr	gga Gly 905	gag Glu		2859
agt Ser	acc Thr	ttt Phe	ttg Leu 910	gta Val	gaa Glu	tgc Cys	act Thr	gag Glu 915	aca Thr	gcg Ala	tca Ser	gtt Val	ctt Leu 920	cag Gln	aat Asn		2907
gca Ala	act Thr	cag Gln 925	gat Asp	tca Ser	cta Leu	gta Val	atc Ile 930	ctt Leu	gac Asp	gaa Glu	ctg Leu	ggc Gly 935	aga Arg	gga Gly	act Thr		2955
agt Ser	act Thr 940	ttc Phe	gat Asp	gga Gly	rac Tyr	gcc Ala 945	acc Ile	gca Ala	tac Tyr	tcg Ser	gtt Val 950	ttt Phe	cgt Arg	cac His	ctg Leu		3003
gta Val 955	gag Glu	aaa Lys	gtt Val	caa Gln	tgt Cys 960	cgg Arg	atg Met	ctc Leu	ttt Phe	gca Ala 965	aca Thr	cat His	tac Tyr	cac His	cct Pro 970		3051
ctc Leu	acc Thr	aag Lys	gaa Glu	ttc Phe 975	gcg Ala	tct Ser	cac His	cca Pro	cgt Arg 980	gtc Val	acc Thr	tcg Ser	aaa Lys	cac His 985	atg Met		3099
gct Ala	tgc Cys	gca Ala	ttc Phe 990	aaa Lys	tca Ser	aga Arg	tct Ser	gat Asp 995	tat Tyr	caa Gln	cca Pro	Arg	ggt Gly 1000	tgt Cys	gat Asp		3147
caa Gln	Asp	cta Leu 1005	gtg Val	ttc Phe	ttg Leu	Tyr	cgt Arg 1010	tta Leu	acc Thr	gag Glu	Gly	gct Ala 015	tgt Cys	cct Pro	gag Glu	;	3195
Ser	tac Tyr 1020	gga Gly	ctt Leu	caa Gln	Val	gca Ala L025	ctc Leu	atg Met	gct Ala	Gly	ata Ile L030	cca Pro	aac Asn	caa Gln	gtg Val	;	3243
gtt Val 103	Glu	aca Thr	gca Ala	Ser	ggt Gly L040	gct Ala	gct Ala	caa Gln	Ala	atg Met 1045	aag Lys	aga Arg	tca Ser	att Ile	050 Gly ggg	;	3291

gga Glu	aac Asn	ttc Phe	Lys	tca Ser .055	agt Ser	gag Glu	cta Leu	Arg	tct Ser .060	gag Glu	ttc Phe	tca Ser	Ser	ctg Leu 1065	cat His	3339
gaa Glu	gac Asp	tgg Trp 1	ctc Leu .070	aag Lys	tca Ser	t tg Leu	Val	ggt Gly L075	att Ile	tct Ser	cga Arg	Val	gcc Ala .080	cac His	aac Asn	3387
aat Asn	Ala	ccc Pro .085	att Ile	ggc Gly	gaa Glu	Asp	gac Asp 1090	tac Tyr	gac Asp	act Thr	Leu	ttt Phe .095	tgc Cys	tta Leu	tgg Trp	3435
His		atc Ile			Ser						taaa	tggc	ta			3478
tgad	ataa	aca c	tato	tgaa	ag ct	cgtt	aagt	ctt	ttg	ctc	tctg	gatgt	tt a	attco	ctctta	3538
aaaa	atgo	ett a	tata	ıtcaa	aa aa	atte	gttt	cto	gatt	aaa	aaaa	aaaa	aaa a	aaaa	aaaaaa	3598
aaaa	ıaaaa	ı	-													3606
<210 <211 <212	L >		31 110 PRT													
<213						tha MSH		na ec	otyp	e Co	lumb	oia				
< 400) >		31	•••												
Met 1	Gln	Arg	Gln	Arg 5	Ser	Ile	Leu	Ser	Phe 10	Phe	Gln	Lys	Pro	Thr 15	Ala	
Ala	Thr	Thr	Lys 20	Gly	Leu	Val	Ser	Gly 25	Asp	Ala	Ala	Ser	Gly 30	Gly	Gly	
Gly	Ser	Gly 35	Gly	Pro	Arg	Phe	Asn 40	Val	Arg	Glu	Gly	Asp 45	Ala	Lys	Gly	
Asp	Ala 50	Ser	Val	Arg	Phe	Ala 55	Val	Ser	Lys	Ser	Val 60	Asp	Glu	Val	Arg	
Gly 65	Thr	Asp	Thr	Pro	Pro 70	Glu	Lys	Val	Pro	Arg 75	Arg	Val	Leu	Pro	Ser 80	
Gly	Phe	Lys	Pro	Ala 85	Glu	Ser	Ala	Gly	Asp 90	Ala	Ser	Ser	Leu	Phe 95	Ser	
Asn	Ile	Met	His 100	Lys	Phe	Val	Lys	Val 105	Asp	Asp	Arg	Asp	Cys 110	Ser	Gly	
Glu	Arg	Ser 115	Arg	Glu	Asp	Val	Val 120	Pro	Leu	Asn	Asp	Ser 125	Ser	Leu	Cys	

	-		
			•
			•

Met	Lys 130	Ala	Asn	Asp	Val	Ile 135	Pro	Gln	Phe	Arg	Ser 140	Asn	Asn	Gly	Lys
Thr 145	Gln	Glu	Arg	Asn	His 150	Ala	Phe	Ser	Phe	Ser 155	Gly	Arg	Ala	Glu	Leu 160
Arg	Ser	Val	Glu	Asp 165	Ile	Gly	Val	Asp	Gly 170	Asp	Val	Pro	Gly	Pro 175	Glu
Thr	Pro	Gly	Met 180	Arg	Pro	Arg	Ala	Ser 185	Arg	Leu	Lys	Arg	Val 190	Leu	Glu
Asp	Glu	Met 195	Thr	Phe	Lys	Glu	Asp 200	Lys	Val	Pro	Val	Leu 205	Asp	Ser	Asn
Lys	Arg 210	Leu	Lys	Met	Leu	Gln 215	Asp	Pro	Val	Cys	Gly 220	Glu	Lys	Lys	Glu
Val 225	Asn	Glu	Gly	Thr	Lys 230	Phe	Glu	Trp	Leu	Glu 235	Ser	Ser	Arg	Ile	Arg 240
qzA	Ala	Asn	Arg	Arg 245	Arg	Pro	Asp	Asp	Pro 250	Leu	Tyr	Asp	Arg	Lys 255	Thr
Leu	His	Ile	Pro 260	Pro	Asp	Val	Phe	Lys 265	Lys	Met	Ser	Ala	Ser 270	Gln	Lys
Gln	Tyr	Trp 275	Ser	Val	Lys	Ser	Glu 280	Tyr	Met	Asp	Ile	Val 285	Leu	Phe	Phe
Lys	Val 290	Gly	Lys	Phe	Tyr	Glu 295	Leu	Tyr	Glu	Leu	Asp 300	Ala	Glu	Leu	Gly
His 305	Lys	Glu	Leu	Asp	Trp 310	Lys	Met	Thr	Met	Ser 315	Gly	Val	Gly	Lys	Cys 320
Arg	Gln	Val	Gly	Ile 325	Ser	Glu	Ser	Gly	Ile 330	_	Glu	Ala	Val	Gln 335	Lys
Leu	Leu	Ala	Arg 340	Gly	Tyr	Lys	Val	Gly 345	Arg	Ile	Glu	Gln	Leu 350	Glu	Thr
Ser	Asp	Gln 355		Lys	Ala	Arg	Gly 360	Ala	Asn	Thr	Ile	Ile 365	Pro	Arg	Lys
Leu	Val 370	Gln	Val	Leu	Thr	Pro 375		Thr	Ala	Ser	Glu 380		Asn	Ile	Gly
Pro 385	Asp	Ala	Val	His	Leu 390	Leu	Ala	Ile	Lys	395	Ile	Lys	Met	Glu	Leu 400
Gln	Lys	Cys	Ser	Thr	Val	Tyr	Gly	Phe	Ala 410		val	Asp	Cys	Ala 415	

			·.	
	÷.			
				·
				•

Leu	Arg	Phe	Trp 420	Val	Gly	Ser	Ile	Ser 425	Asp	Asp	Ala	Ser	Cys 430	Ala	Ala
Leu	Gly	Ala 435	Leu	Leu	Met	Gln	Val 440	Ser	Pro	Lys	Glu	Val 445	Leu	Tyr	Asp
Ser	Lys 450	Gly	Leu	Ser	Arg	Glu 455	Ala	Gln	Lys	Ala	Leu 460	Arg	Lys	Tyr	Thr
Leu 465	Thr	Gly	Ser	Thr	Ala 470	Val	Gln	Leu	Ala	Pro 475	Val	Pro	Gln	Val	Met 480
Gly	qzA	Thr	Asp	Ala 485	Ala	Gly	Val	Arg	Asn 490	Ile	Ile	Glu	Ser	Asn 495	Gly
Tyr	Phe	Lys	Gly 500	Ser	Ser	Glu	Ser	Trp 505	Asn	Cys	Ala	Val	Asp 510	Gly	Leu
Asn	Glu	Cys 515	Asp	Val	Ala	Leu	Ser 520	Ala	Leu	Gly	Glu	Leu 525	Ile	Asn	His
Leu	Ser 530	Arg	Leu	Lys	Leu	Glu 535	Asp	Val	Leu	Lys	His 540	Gly	Asp	Ile	Phe
Pro 545	Tyr	Gln	Val	туг	Arg 550	Gly	Cys	Leu	Arg	Ile 555	Asp	Gly	Gln	Thr	Met 560
Val	Asn	Leu	Glu	Ile 565	Phe	Asn	Asn	Ser	Cys 570	Asp	Gly	Gly	Pro	Ser 575	Gly
Thr	Leu	Tyr	Lys 580	Tyr	Leu	Asp	Asn	Cys 585	Val	Ser	Pro	Thr	Gly 590	Lys	Arg
Leu	Leu	Arg 595	Asn	Trp	Ile	Cys	His 600	Pro	Leu	Lys	Asp	Val 605	Glu	Ser	Ile
Asn	Lys 610	Arg	Leu	Asp	Val	Val 615	Glu	Glu	Phe	Thr	Ala 620	Asn	Ser	Glu	Ser
Met 625	Gln	Ile	Thr	Gly	Gln 630	Tyr	Leu	His	Lys	Leu 635	Pro	Asp	Leu	Glu	Arg 640
Leu	Leu	Gly	Arg	Ile 645		Ser	Ser	Val	Arg 650		Ser	Ala	Ser	Val 655	Leu
Pro	Ala	Leu	Leu 660		Lys	Lys	Val	Leu 665		Gln	Arg	Val	Lys 670	Ala	Ph∈
Gly	Gln	Ile 675		Lys	Gly	Phe	Arg 680		Gly	'Ile	Asp	Leu 685		Leu	Ala
Leu	Gln		Glu			Met		Ser	Leu	Leu	Tyr		Leu	Cys	Lys

			•	
	*			

Leu 705	Pro	Ile	Leu	Val	Gly 710	Lys	Ser	Gly	Leu	Glu 715	Leu	Phe	Leu	Ser	Gln 720
Phe	Glu	Ala	Ala	Ile 725	Asp	Ser	Asp	Phe	Pro 730	Asn	Tyr	Gln	Asn	Gln 735	Asp
Val	Thr	Asp	Glu 740	Asn	Ala	Glu	Thr	Leu 745	Thr	Ile	Leu	Ile,	Glu 750	Leu.	Phe
Ile	Glu	Arg 755	Ala	Thr	Gln	Trp	Ser 760	Glu	Val	Ile	His	Thr 765	Ile	Ser	Cys
Leu	Asp 770	Val	Leu	Arg	Ser	Phe 775	Ala	Ile	Ala	Ala	Ser 780	Leu	Ser	Ala	Gly
Ser 785	Met	Ala	Arg	Pro	Val 790	Ile	Phe	Pro	Glu	Ser 795	Glu	Ala	Thr	Asp	Gln 800
Asn	Gln	Lys	Thr	Lys 805	Gly	Pro	Ile	Leu	Lys 810	Ile	Gln	Gly	Leu	Trp 815	His
Pro	Phe	Ala	Val 820	Ala	Ala	qsA	Gly	Gln 825	Leu	Pro	Val	Pro	Asn 830	Asp	Ile
Leu	Leu	Gly 835	Glu	Ala	Arg	Arg	Ser 840	Ser	Gly	Ser	Ile	His 845	Pro	Arg	Ser
Leu	Leu 850	Leu	Thr	Gly	Pro	Asn 855	Met	Gly	Gly	Lys	Ser 860	Thr	Leu	Leu	Arg
Ala 865	Thr	Cys	Leu	Ala	Val 870	Ile	Phe	Ala	Gln	Leu 875	Gly	Cys	Tyr	Val	Pro 880
Cys	Glu	Ser	Cys	Glu 885	Ile	Ser	Leu	Val	Asp 890	Thr	Ile	Phe	Thr	Arg 895	Leu
Gly	Ala	Ser	Asp 900	Arg	Ile	Met	Thr	Gly 905	Glu	Ser	Thr	Phe	Leu 910	Val	Glu
Cys	Thr	Glu 915	Thr	Ala	Ser	Val	Leu 920	Gln	Asn	Ala	Thr	Gln 925	Asp	Ser	Leu
Val	Ile 930	Leu	Asp	Glu	Leu	Gly 935	Arg	Gly	Thr	Ser	Thr 940	Phe	Asp	Gly	Tyr
Ala 945	Ile	Ala	Tyr	Ser	Val 950	Phe	Arg	His	Leu	Val 955	Glu	Lys	Val	Gln	Cys 960
Arg	Met	Leu	Phe	Ala 965	Thr	His	Tyr	His	Pro 970	Leu	Thr	Lys	Glu	Phe 975	Ala
Ser	His	Pro	Arg 980	Val	Thr	Ser	Lys	His 985	Met	Ala	Cys	Ala	Phe 990	Lys	Ser

,		,		•	
	4.				
					•
					•

WO 99/19492 PCT/EP98/06977

Arg Ser Asp 995	Tyr Gln Pro Arg Gly Cys Asp Gln Asp Leu Val Phe Leu 1000 1005	
Tyr Arg Leu 1010	Thr Glu Gly Ala Cys Pro Glu Ser Tyr Gly Leu Gln Val 1015 1020	
Ala Leu Met 1025	Ala Gly Ile Pro Asn Gln Val Val Glu Thr Ala Ser Gly 1030 1035 1040	
Ala Ala Gln	Ala Met Lys Arg Ser Ile Gly Glu Asn Phe Lys Ser Ser 1045 1050 1055	
	Ser Glu Phe Ser Ser Leu His Glu Asp Trp Leu Lys Ser 1060 1065 1070	
Leu Val Gly 1075	Ile Ser Arg Val Ala His Asn Asn Ala Pro Ile Gly Glu 1080 1085	
Asp Asp Tyr 1090	Asp Thr Leu Phe Cys Leu Trp His Glu Ile Lys Ser Ser 1095 1100	
Tyr Cys Val 1105	Pro Lys	
<210> <211> <212> <213>	32 24 DNA Artificial sequence	
<220> <223>	Forward primer for PCR amplification of ATHGENEA microsatellite	
<400>	32	
accatgcata	gcttaaactt cttg	24
<210> <211> <212> <213>	33 22 DNA Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of ATHGENEA microsatellite	
<400>	33	
acataaccac	aaataggggt gc	22

	,		·	
·		`` '\		·
		"		•
	÷		÷	

WO 99/19492 PCT/EP98/06977

<210>	34				
<211>	18				
<212>	DNA				
<213>	Artificial sequence				
<220>					
<223>	Forward primer DMCIN-A for PCR thaliana ssp. Landsberg erecta	on genomic "Ler"	DNA	of	Arabidopsis
<400>	34				
gaagcgatat	tgttcgtg				18
<210>	35				
<211>	18				
<212>	DNA				
<213>	Artificial sequence				
<220>					
<223>	Reverse primer DMCIN-B for PCR thaliana ssp. Landsberg erecta	on genomic "Ler"	DNA	of	Arabidopsis
<400>	35				
agat tgcgag	aacattcc				18
<210>	36				
<211>	31				
<212>	DNA				
<213>	Artificial sequence				
<220>					
<223>	Forward primer DMCIN-1 for PCR thaliana ssp. Landsberg erecta	on genomic "Ler"	DNA	of	Arabidopsis
<400>	36				
acgcgtcgac	tcagctatga gattactcgt g				31
<210>	37				
<211>	29				
<212>	DNA				
<213>	Artificial sequence				
<220>					
<223>	Reverse primer DMCIN-2 for PCR thaliana ssp. Landsberg erecta	on genomic "Ler"	DNA	of	Arabidopsis
<400>	37				
gctctagatt	totogotota agactotot				29

			<i>*</i>	
	<u>.</u>			
				•

<212>	38 32 DNA Artificial sequence
<220 > <223 >	Forward primer DMCIN-3 for PCR on genomic DNA of Arabidopsis thaliana ssp. Landsberg erecta "Ler"
<400>	38
gctctagage t	tctcttaag taagtgattg at 32
<211> <212>	39 48 DNA Artificial sequence
<220>	
<223>	Reverse primer DMCIN-4 for PCR on genomic DNA of Arabidopsis thaliana ssp. Landsberg erecta "Ler"
<400>	39
teeceegge t	cgagagatc tocatggttt cttcagctct atgaatcc 48
<210> <211> <212> <213>	40 26 DNA Artificial sequence
<220> <223>	Forward primer DMCla for PCR on genomic DNA of Arabidopsis thaliana ssp. Landsberg erecta "Ler"
<400>	40
acgcgtcgac g	aattcgcaa gtgggg 26
<210> <211> <212> <213>	41 38 DNA Artificial sequence
<220> <223>	Reverse primer DMC1b for PCR on genomic DNA of Arabidopsis thaliana ssp. Landsberg erecta "Ler"
<400>	41

			· 3.	·	
	P. g				
				•	

WO 99/19492 PCT/EP98/06977

	→	
tccatggaga	totocogggt acogatttgo ttogaggg	38
<210>	42	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of ATEAT1 SSLP mar Arabidopsis thaliana subspecies	rker in
<400>	42	
g ccactg egt	gaatgatatg	20
<210>	43	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of ATEAT1 SSLP man Arabidopsis thaliana subspecies	rker in
<400>	43	
cgaacagcca	acattaattc cc	22
.210		
<210><211>	44	
<211>	18	
<213>	DNA Artificial sequence	
(213)	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA63 SSLP mark Arabidopsis thaliana subspecies	ker in
<400>	44	
aaccaaggca	cagaagcg	18
<210>	45	
<211>	18	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA63 SSLP mar. Arabidopsis thaliana subspecies	ker in

	÷.			
			"	•
			19	
•				

NO 99/19492	PCŢ/EP98/06977

	36	
	•	
< 400>	45	
acccaagtga	tegecace	18
<210>	46	
<211>	21	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA248 SSLP Arabidopsis thaliana subspecies	marker in
<400>	46	
taccgaacca	aaacacaaag g	21
<210>	47	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA248 SSLP Arabidopsis thaliana subspecies	marker in
<400>	47	
tctgtatctc	ggtgaattet ee	22
<210>	48	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA128 SSLP	marker in
12237	Arabidopsis thaliana subspecies	marker II
<400>	48	
ggtctgttga	tgtcgtaagt cg	22
<210>	40	
	49	
<211>	22	
<212> <213>	DNA	
(213)	Artificial sequence	
<220>		

				<u>;</u>	
		æ,			
	190				
i.					
			el.		

<223>	Reverse primer for PCR amplification of NGA128 SSLP marker in Arabidopsis thaliana subspecies
<400>	49
atcttgaaac c	tttagggag gg 22
<210>	50
<211>	22
<212>	DNA
<213>	Artificial sequence
(213)	Attitital sequence
<220>	
<223>	Forward primer for PCR amplification of NGA280 SSLP marker in Arabidopsis thaliana subspecies
<400>	50
ctgatctcac g	gacaatagt gc 22
<210>	51
<211>	20
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of NGA280 SSLP marker in Arabidopsis thaliana subspecies
< 400 >	51
ggctccataa a	aaagtgcacc 20
<210>	52
<211>	21
<212>	DNA
<213>	
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA111 SSLP marker in
	Arabidopsis thaliana subspecies
<400>	52
ctccagttgg a	aagctaaagg g 21
<210>	53
<211>	21
<211>	DNA
<212> <213>	Artificial sequence
\~ L J >	wrettretat acdresses

			-10	
2 20				
		•		

PCŢ/EP98/06977

WO 99/19492

cgctacgctt ttcggtaaag

	•				
<220>					
< 223 >	Reverse primer for PCR amplification	of NGA111	SSLP	marker	in
	Arabidopsis thaliana subspecies				
	•				
<400>	53				
(400)	55				
tgttttttag g	acaaatgge g	*	-		21
		**			
<210>	54				
<211>	20				
<212>	DNA				
<213>	Artificial sequence				
	-				
<220>					
<223>	Forward primer for PCR amplification	of NGA168	SSLP	marker	in
(225)	Arabidopsis thaliana subspecies	OI MONTOO	2011	MULTICE	
	Alabidopsis challana subspecies				
400					
< 400>	54				
cetteacate e	aaaacccac				20
<210>	55				
<211>	20				
<212>	DNA				
<213>	Artificial sequence				
	The call of a control				
<220>					
<223>	Borowca primar for DCB amplification	of NCA169	CCID	marker	· in
(223)	Reverse primer for PCR amplification	OI NGA100	2275	marker	T 1 1
	Arabidopsis thaliana subspecies				
<400>	55				
gcacataccc a	acaaccagaa				20
<210>	56				
<211>	20				
<212>	DNA				
<213>	Artificial sequence				
-417	Wrettrerar poducines				
-220>					
<220>		,		5 •	
<223>	Forward primer for PCR amplification	of NGA112	ь SSI	r marke	E
	in Arabidopsis thaliana subspecies				
<400>	56				
	•				

38

,	,		•	

PCT/EP98/06977

39

WO 99/19492

-	•
<210>	57
<211>	20
	DNA
<213>	Artificial sequence
	•
<220>	
<223>	Reverse primer for PCR amplification of NGA1126 SSLP marker
	in Arabidopsis thaliana subspecies
< 400>	57
gcacagteca a	gtcacaacc 20
<210>	58
<211>	20
<212>	DNA
<213>	Artificial sequence
(213)	metretur begaenee
<220>	
<223>	Forward primer for PCR amplification of NGA361 SSLP marker in
	Arabidopsis thaliana subspecies
<400>	58
aaagagatga g	gaatttggac 20
<210>	59
<211>	23
<212>	DNA
<213>	Artificial sequence
	1,000
<220>	
<223>	Reverse primer for PCR amplification of NGA361 SSLP marker in
	Arabidopsis thaliana subspecies
<400>	59
acatatcaat a	atattaaagt agc 23
<210>	60
<211>	18
<212>	DNA
<213>	Artificial sequence
	······································
<220>	
<223>	Forward primer for PCR amplification of NGA168 SSLP marker in
	Arabidopsis thaliana subspecies
	•
<400>	60

			•

tegtetactg	cactgoog	18
<210>	· 61	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
(613)	Altilital sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA168 SSLP marker Arabidopsis thaliana subspecies	in
<400>	61	
gaggacatgt	ataggagcct cg	22
<210>	62	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of AthBIO2 SSLP marke in Arabidopsis thaliana subspecies	er
<400>	62	
tgacctcctc	ttccatggag	20
<210>	63	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of AthBIO2 SSLP marks in Arabidopsis thaliana subspecies	er
<400>	63	
ttaacagaaa	cccaaagett te	22
<210>	64	
<211>	21	
<212>	DNA	
<213>		
	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of AthUBIQUE SSLP main Arabidopsis thaliana subspecies	rker

		÷	
4			
			•

```
<400>
             64
aggcaaatgt ccatttcatt g
                                                                         21
<210>
             65
<211>
             20
<212>
             DNA
<213>
             Artificial sequence
<220>
             Reverse primer for PCR amplification of AthUBIQUE SSLP marker
<223>
             in Arabidopsis thaliana subspecies
<400>
             65
                                                                         20
acgacatggc agatttctcc
<210>
             66
<211>
             21
<212>
             DNA
<213>
             Artificial sequence
<220>
<223>
             Forward primer for PCR amplification of NGA172 SSLP marker in
             Arabidopsis thaliana subspecies
<400>
             66
agctgcttcc ttatagcgtc c
                                                                         21
<210>
             67
<211>
             19
<212>
             DNA
<213>
             Artificial sequence
<220>
<223>
             Reverse primer for PCR amplification of NGA172 SSLP marker in
             Arabidopsis thaliana subspecies
<400>
             67
catccgaatg ccattgttc
                                                                         19
<210>
             68
<211>
              21
<212>
             DNA
<213>
             Artificial sequence
<220>
```

	,			
		9		
4.7				
				٠
				•

WO 99/19492

< 323 >	Forward primer for PCR amplification of NGA126 SSLP marker in Arabidopsis thaliana subspecies
< 400>	63
gaaaaaacgc	tactttcgtg g 21
<210>	69
<211>	22
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of NGA126 SSLP marker in Arabidopsis thaliana subspecies
<400>	69
caagagcaat	atcaagagca gc 22
<210>	70 -
<211>	20
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA162 SSLP marker in Arabidopsis thaliana subspecies
<400>	70
catgcaattt	gcatctgagg 20
<210>	71
<211>	22
<212>	DNA
<213>	Artificial sequence
(213)	Arctificial sequence
<220>	•
<223>	Reverse primer for PCR amplification of NGA162 SSLP marker in Arabidopsis thaliana subspecies
<400>	71
ctctgtcact	cttttectct gg 22
<210>	72
<211>	72 21
<212>	DNA
<213>	Artificial sequence

					• =	
		\ -				•
				**		
						•
	•					
		•				
-5						
			190			

PCŢ/EP98/06977

<220>		
<223>	Forward primer for PCR amplification of NGA6 SSLP marker is Arabidopsis thaliana subspecies	n
< 400 >	72	
tggatttett e	ctctcttca c	21
<210>	73	
<211>	21	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA6 SSLP marker i Arabidopsis thaliana subspecies	.n
<400>	73	
atggagaagc t	tacactgat c	21
<210>	74	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA12 SSLP marker Arabidopsis thaliana subspecies	in
<400>	74	
aatgttgtcc t	cccctcctc	20
210		
<210>	75	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA12 SSLP marker Arabidopsis thaliana subspecies	in
<400>	75	
tgatgctctc t	tgaaacaaga gc	22

22

		-9,	
•			
**			
			•
			,
			2)

PCT/EP98/06977

<210>	76
<211>	21
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA8 SSLP marker in Arabidopsis thaliana subspecies
<400>	76
gagggcaaat	ctttatttcg g 21
<210>	77
<211>	22
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of NGA8 SSLP marker in Arabidopsis thaliana subspecies
<400>	77
tggctttcgt	ttataaacat cc 22
<210>	78
<211>	21
<212>	DNA
<213>	
(213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA1107 SSLP marker in Arabidopsis thaliana subspecies
<400>	78
gcgaaaaaac	aaaaaaatcc a 21
ı	_
<210>	79
<211>	21
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of NGA1107 SSLP marker in Arabidopsis thaliana subspecies
<400>	79

			4.	

cgacgaatcg	acagaattag _g	21
<210>	. 80	
<211>	21	
<212>	DNA	
<213>	Artificial sequence	
<220>	•	
<223>	Forward primer for PCR amplification of NGA225 SSLP marke: Arabidopsis thaliana subspecies	r in
<400>	80	
gaaatccaaa	tcccagagag g	21
<210>	81	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA225 SSLP marke. Arabidopsis thaliana subspecies	r in
<400>	81	
tctccccact	agttttgtgt cc	22
210		
<210>	82	
<211>	19	
<212> <213>	DNA Artificial sequence	
	Altilitat soquence	
<220>		
<223>	Forward primer for PCR amplification of NGA249 SSLP marke Arabidopsis thaliana subspecies	r in
<400>	82	
taccgtcaat	ttcatcgcc	19
<210>	83	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
· -		
<220>	·	
<223>	Reverse primer for PCR amplification of NGA249 SSLP marke Arabidopsis thaliana subspecies	r in

				, ,	
	4.		,		
j.			"		<u> </u>
		,			

<400> -	. 83
ggateectaa	ctgtaaaatc cc 22
<210>	84 -
<211>	22
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of CA72 SSLP marker in Arabidopsis thaliana subspecies
<400>	84
aatcccagta	accaaacaca ca 22
<210>	85
<211>	20
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of CA72 SSLP marker in Arabidopsis thaliana subspecies
<400>	85
cccagtctaa	ccacgaccac 20
<210>	86
<211>	20
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA151 SSLP marker in Arabidopsis thaliana subspecies
<400>	86
gttttgggaa	a gttttgctgg 20
<210>	87
<211>	24
<212>	DNA
<213>	Artificial sequence
<220>	

	÷,		
			-

WO 99/19492 PCT/EP98/06977

47

<223>	Reverse primer for PCR amplification of NGA151 SSLP marker i Arabidopsis thaliana subspecies	n
<400>	87	
cagtctaaaa	gcgagagtat gatg	t
<210>	88	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA106 SSLP marker i Arabidopsis thaliana subspecies	n
<400>	88	
gttatggagt	ttctagggca cg 22	į
<210>	89	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
(213)	Arciriciai sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA106 SSLP marker i Arabidopsis thaliana subspecies	.n
< 400>	89	
tgccccattt	tgttcttctc 20)
<210>	90	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primar for DCD "amplification of worden corp.	_
	Forward primer for PCR amplification of NGA139 SSLP marker i Arabidopsis thaliana subspecies	.n
<400>	90	
agagctacca	gatecgatgg 20)
<210>	91	
<211>	21	
<212>	DNA	
<213>	Artificial sequence	
	· · · · · · · · · · · · · · · · · · ·	

			,	÷	
NA S	-				32
					•

WO 99/19492 PCT/EP98/06977

48

<220>						
<223>	Reverse primer for PCR amplification Arabidopsis thaliana subspecies	of	NGA139	SSLP	marker	in
<400>	91					
ggtttcgttt c	actatccag g	, ,,				21
<210>	92					
<211>	22					
<212>	DNA					
<213>	Artificial sequence					
<220>						
<223>	Forward primer for PCR amplification Arabidopsis thaliana subspecies	of	NGA76	SSLP	marker	in
<400>	92					
ggagaaaatg t	cacteteca ec					22
<210>	93					
<211>	20					
<212>	DNA					
<213>	Artificial sequence					
<220>						
<223>	Reverse primer for PCR amplification Arabidopsis thaliana subspecies	of	NGA76	SSLP	marker	in
<400>	93					
aggcatggga g	acatttacg					20
<210>	94					
<211>	20					
<212>	DNA					
<213>	Artificial sequence					
- was with her P	merracial seducines					
<220>						
<223>	Forward primer for PCR amplification in Arabidopsis thaliana subspecies	of	ATHSO1	91 SS	SLP mark	er
<400>	94					
ctccaccaat c	atgcaaatg					20

•			
			Ą
			-

•	•
<210>	95
	21
<211>	
<212>	· DNA
<213>	Artificial sequence
	-
<220>	5 ATMOST AL CONTRACTOR AND ANALOGO ANALOGO AND ANALOGO ANA
<223>	Reverse primer for PCR amplification of ATHSO191 SSLP marker
	in Arabidopsis thaliana subspecies
	-
<400>	95
tgatgttgat g	gagatggtc a 21
, , , , ,	,
<210>	96
<211>	22
<212>	DNA
<213>	Artificial sequence
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA129 SSLP marker in
,	Arabidopsis thaliana subspecies
	Arabidopsis charrana Sabbectes
<400>	96
traggaggaa	ctaaagtgag gg 22
0003303300	
<210>	97
<211>	22
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of NGA129 SSLP marker in
12237	Arabidopsis thaliana subspecies
	Arabidopsis tharrana subspecies
<400>	97
	•
and a crange	atggtcttga gg 22
cacactyaay	atggtcttga gg 22
	-
<210>	98
<211>	8062
<212>	DNA
<213>	Arabidopsis thaliana ecotype Columbia
<220>	
	Conomic DNA sequence of AtMSUE
<223>	Genomic DNA sequence of AtMSH6
<400>	97

		*	Ģ-
- 4			
			4

aagaaatgaa agatatatat tgttttttca tttatcaaac aaaacaacaa gactttttt 120 180 ttacttttta cattggtcaa caaaatacaa gataaacgac atcgtttaat catttcccaa ttttacccct aagtttaaca cctagaacct tctccatctt cgcaagcaca gcctgattag 240 gaacagettt accattetea tatteetgaa etaeetgagt eeteteattg atetgttteg 300 360 ccaaatccgc ttgtgacatc ttcttctcca atctcgcttt ctgtatcatc aacctcacct ctgctttcac acgatccatc gccgcaggct ctgtttcttc ttccagcttc ttcgtgttaa 420 tcaccggaac cgccgtagat ttcccctttt tgttcgaacc ggcatcgaat ttcttaaccg 480 tttgaaccgc gacaccgttt ctcagagctg cgttaaccgc tttcggatcg cgtaggtctt 540 600 ggetettttg ttttgatttg tggagaacta etggtteeca gtettgtgtt aetgeteetg 660 qqtatctqct cggcatcgtc gatgaattga gagaaaggaa caacgcgaaa attttattaa 720 totgagtttt gaaattgaga aacgatgaag atgaagaatg ttgttgagag gattgtgata 780 tttatatata egaagattgg tttetggaga attegateat ettttetee attttegtet 840 ctggaacgtt cttagagatg attgacgacg tgtcattatc tgatttgcag ttaaccaatg 900 ctttttgggt tggattegtg gtacaccata ttateegatt tggeteaatg gttttatata 960 aatttggttt teggtteggt tatgagttat eattaaaatt aagetaacea aaaatttteg 1020 taaaatttat tteggtttea atteggatee ettaetteea gaacegaatt attegaaace 1080 ggggttagee gaacegaata ecaatgeetg attgactegt tggetagaaa gateeaacgg 1140 tatacaataa tagaacataa atcggacggt catcaaagcc tcaaagagtg aacagtcaac 1200 aaaaaaagtt gagccctgag gagtatcgtt tccgccattt ctacgacgca aggcgaaaat ttttggcgcc aatotttccc ccctttcgaa ttctctcagc tcaaaacatc gtttctctct 1260 1320 cactetetet cacaatteca aaaaatgeag egecagagat egattttgte tttetteeaa 1380 aaacccacgg cggcgactac gaagggtttg gtttccggcg atgctgctag cggcgggggc 1440 ggcagcggag accacgattt aatgtgaagg aaggggatge taaaggcgac gettetgtac 1500 gttttgctgt ttcgaaatct gtcgatgagg ttagaggaac ggatactcca ccggagaagg ttccgcgtcg tgtcctgccg tctggattta agccggctga atccgccggt gatgcttcgt 1560 1620 ccctgttctc caatattatg cataagtttg taaaagtcga tgatcgagat tgttctggag 1680 agaggtacta atottogatt ctottaattt tgttatottt agotggaaga agaagattog

					1.
				•	
		•			
	- •				
					4
					•
			-		
•					

WO 99/19492 PCT/EP98/06977

tgtaatttgt	tgtattcgtt	ggagagattc	tgattactgc	attggatcgt	tgtttacaaa	1740
ttttcaggag	ccgagaagat	gttgttccgc	tgaatgattc	atctctatgt	atgaaggcta	1800
atgatgttat	tcctcaattt	cgttccaata	atggtaaaac	tcaagaaaga	aaccatgctt	1860
ttagtttcag	tgggagagct	gaacttagat	cagtagaaga	tataggagta	gatggcgatg	1920
ttcctggtcc	agaaacacca	gggatgcgtc	cacgtgcttc	tcgcttgaag	cgagttctgg	1980
aggatgaaat	gacttttaag	gaggataagg	ttcctgtatt	ggactctaac	aaaaggctga	2040
aaatgctcca	ggatccggtt	tgtggagaga	agaaagaagt	aaacgaagga	accaaatttg	2100
aatggcttga	gtcttctcga	atcagggatg	ccaatagaag	acgtcctgat	gateceettt	2160
acgatagaaa	gaccttacac	ataccacctg	atgttttcaa	gaaaatgtct	gcatcacaaa	2220
agcaatattg	gagtgttaag	agtgaatata	tggacattgt	gcttttcttt	aaagtggtta	2280
gtaactatta	atctagtgtt	caatccattt	cctcaatgtg	atttgttcac	ttacatctgt	2340
ttacgttatg	ctcttctcag	gggaaatttt	atgagctgta	tgagctagat	gcggaattag	2400
gtcacaagga	gcttgactgg	aagatgacca	tgagtggtgt	gggaaaatgc	agacaggtaa	2460
attagttgaa	acaactggcc	tgcttgaatt	attgtgtcta	taaattttga	caccaccttt	2520
tgtttcaggt	tggtatctct	gaaagtggga	tagatgaggc	agtgcaaaag	ctattagete	2580
g tgggtaag g	gaaccatcat	actttatgga	attcgtttac	tgctacttcg	gctaggattt	2640
aagaaatgga	aatcacttca	agcatcatta	gttaggatcc	tgagaactca	ggatgttttc	2700
ttattcgtta	tataataagt	cttttcatca	aggagtaaca	aacaaaactt	gcacaatatt	2760
tgtgtgctca	ctggcaaggc	atatataccc	agctaacctt	tgctagttca	ctgtagtaac	2820
agttacggat	aatatatgtt	tacttgtatg	tggtaccctc	attttgtctc	tcatggaggc	2880
tttcaagcct	tgtgttgaaa	ctggatagtt	acatatgctt	ccaacagaaa	ctagcatgca	2940
gattcatatg	ctttcctatt	ctactaatta	tgtattgaca	cactcgttgt	ttcttttgaa	3000
agatataaag	ttggacgaat	cgagcagcta	gaaacatctg	accaagcaaa	agccagaggt	3060
gctaatactg	taagttttct	tggataggtc	aaggagagtg	ttgcagactg	tttttgatca	3120
tttcttttc	tgtacattac	tttcatgctg	taattaactc	aatggctatt	ctggtctgat	3180
tatcagataa	ttccaaggaa	gctagttcag	gtattaactc	catcaacagc	aagcgaggga	3240
aacatcgggc	ctgatgccgt	ccatcttctt	gctataaaag	aggtttgtta	tttacttatt	3300

	-	,	•	
•			1. - 1	•

•	•	-				
tatottatoa	tgttcagttc	atccaagtcc	tgaaaaatta	cactettett	taccaatctt	3360
ccatcaagct	gtgtaaagga	tttggaatta	gaaaatcatt	atttgatgct	ttgttttata	3420
tgcaagaggt	tcccttgaaa	agatctgttt	aagattcttt	gcacttgaaa	aattcaatct	3480
ttttaagtga	atcccctact	ttcttacaat	gatcatagtc	tgcaattgca	tgtcaagtaa	3540
tatcattcct	tgttactgca	teccectett	tcttaatgac	cattgtctat	gttgtgtttg	3600
tetegtgtge	tggagaaaat	gatagctgat	ccaagctgta	cattatcatg	attaagtagc	3660
tgctcaggaa	ttgcctttgg	ttacattgcc	taatggtttg	atgtcaattt	ttcttctgaa	3720
tctttattt	agatcaaaat	ggagctacaa	aagtgttcaa	ctgtgtatgg	atttgctttt	3780
gttgactgtg	ctgccttgag	gttttgggtt	gggtccatca	gcgatgatgc	atcatgtgct	3840
gctcttggag	cgttattgat	gcaggtaagc	aagtgtattc	tgtatcttat	gtgtaccatg	3900
tgacttcctg	tgcatatatt	tgggttgcag	gaactaattc	tgaatcacca	tttggtatgt	3960
tttttccagg	tttctccaaa	ggaagtgtta	tatgacagta	aaggtaaact	gcttgtatcg	4020
ccagttgttt	tgttaaacag	aatttaaggt	aaatgacact	ggttaattta	aagtgcatac	4080
atgttgaaat	attgcagggc	tatcaagaga	agcacaaaag	gctctaagga	aatatacgtt	4140
gacaggtacc	atttcagtag	gcaagctaac	tgacaattta	accgctcacc	gaatgatagg	4200
tctcttaaac	attgctaatg	tagatgatgt	ttatgtttca	atctaatagg	gtctacggcg	4260
gtacagttgg	ctccagtacc	acaagtaatg	ggggatacag	atgctgctgg	agttagaaat	4320
ataatagaat	ctaacggata	ctttaaaggt	tcttctgaat	catggaactg	tgctgttgat	4380
ggtctaaatg	aatgtgatgt	tgcccttagt	gctcttggag	agctaattaa	tcatctgtct	4440
aggctaaagg	tgtgttggct	tgtttagttt	ttgcttttca	caaattaagc	aaaggaactt	4500
ttcataactt	acagtttcta	tctacttgca	gctagaagat	gtacttaagc	atggggatat	4560
ttttccatac	caagtttaca	ggggttgtct	cagaattgat	ggccagacga	tggtaaatct	4620
tgagatattt	aacaatagct	gtgatggtgg	tectteagge	aagtgcatat	ttcttttttg	4680
ataacttcaa	ctagagggca	gacatagaag	gaaaaattct	aatacttcgt	acggatetee	4740
agtaagtaat	agccgatttt	tgtttaccta	tgtagggacc	ttgtacaaat	atcttgataa	4800
ctgtgttagt	ccaactggta	agegaetett	aaggaattgg	atctgccatc	cactcaaaga	4860
tgtagaaago	atcaataaac	ggcttgatgt	agttgaagaa	ttcacggcaa	actcagaaag	4920

			•	
				,
				·
	*	- 1		

tatgcaaatc actggccagt átctccacaa acttccagac ttagaaagac tgctcggacg 4980 catcaagtct agegttegat catcageete tgtgttgeet getettetgg ggaaaaaagt 5040 gctgaaacaa cgagtaagta tcaatcacaa gttttctgag taatgccttc catgagtagt 5100 ataggactaa aacattacgg gtctagctaa agactgttct ccttctttg caatgtctqq 5160 ttattcatta catttctctt aacttattgc attgcaggtt aaagcatttg ggcaaattgt 5220 gaaagggttc agaagtggaa ttgatctgtt gttggctcta cagaaggaat caaatatgat 5280 gagtttgctt tataaactct gtaaacttcc tatattagta ggaaaaagcg ggctagagtt 5340 atttctttct caattcgaag cagccataga tagcgacttt ccaaattatc aggtgcccat 5400 ctatctttca tactttacaa caaaatgtct gtcactactc aaagcaatgc atatggctta 5460 gateteaact cacaccega ggatectaaa gggatetget tettatteet aatgttettg 5520 gatggtttga tttatttcta acttgaactt attaatcttg taccagaacc aagatgtgac 5580 agatgaaaac getgaaacte teacaataet tategaaett tttategaaa gageaaetea 5640 atggtctgag gtcattcaca ccataagctg cctagatgtc ctgagatctt ttgcaatcgc 5700 agcaagtete tetgetggaa geatggeeag geetgttatt ttteeegaat cagaagetae 5760 agatcagaat cagaaaacaa aagggccaat acttaaaatc caaggactat ggcatccatt 5820 tgcagttgca gccgatggtc aattgcctgt tccgaatgat atactccttg gcgaggctag 5880 aagaagcagt ggcagcattc atcctcggtc attgttactg acgggaccaa acatgggcgg 5940 6000 aaaatcaact cttcttcgtg caacatgtct ggccgttatc tttgcccaag tttgtatact cgttagataa ttactctatt ctttgcaatc agttcttcaa catgaataat aaattctgtt 6060 ttctgtctgc agcttggctg ctacgtgccg tgtgagtctt gcgaaatctc cctcgtggat 6120 actatettea caaggettgg egeatetgat agaateatga caggagagag taagttttgt 6180 totcaaaata coaattooto gaactattta otcagatttt gtotgattgg acaaggtggt 6240 tttgcttttt tttaggtacc tttttggtag aatgcactga gacagcgtca gttcttcaga 6300 atgcaactca ggattcacta gtaatccttg acgaactggg cagaggaact agtactttcg 6360 atggatacge cattgcatae teggtaacet getettetee tteaacttat acttgttgat 6420 caacaaaaac atgcaattca ttttgctgaa acttattgat ttatatcagg tttttcgtca 6480 cctggtagag aaagttcaat gtcggatgct ctttgcaaca cattaccacc ctctcaccaa 6540

			· (i)			
				•	¿*,	
•						
					•	•
					•	•
	. 2					
	- +					
		,				

			••			
ggaattcgcg	tctcacccac	gtgtcacctc	gaaacacatg	gcttgcgcat	tcaaatcaag	6600
atctgattat	caaccacgtg	gttgtgatca	agacctagtg	ttcttgtacc	gtttaaccga	6660
gggagcttgt	cctgagagct	acggacttca	agtggcactc	atggctggaa	taccaaacca	6720
agtggttgaa	acagcatcag	gtgctgctca	agccatgaag	agatcaattg	gggaaaactt	6780
caagtcaagt	gagctaagat	ctgagttctc	aagtctgcat	gaagactggc	tcaagtcatt	6840
ggtgggtatt	tctcgagtcg	cccacaacaa	tgcccccatt	ggcgaagatg	actacgacac	6900
tttgttttgc	ttatggcatg	agatcaaatc	ctcttactgt	gttcccaaat	aaatggctat	6960
gacataacac	tatctgaagc	tcgttaagtc	ttttgcttct	ctgatgttta	ttcctcttaa	7020
aaaatgctta	tatatcaaaa	aattgtttcc	tcgattataa	caagattata	tatgtatctg	7080
tcggtttagc	tatggtatat	aatatatgta	tgttcatgag	attggtcaag	agaaatactc	7140
acaaacagta	tattaagaag	gaaatatgtt	tatgcattaa	tttaagtttc	aagataaact	7200
gcaaataacc	tcgactaaag	ttgcaaagac	caaacacaaa	ttacaaaact	tataagactt	7260
aagttctgaa	ttccctaaaa	ccaaaaaaaa	aaacagaaca	tattttgttg	catctacaaa	7320
caacacaaac	ctacatagtt	tataacttac	tcatcactga	gattaacatc	agaatcattc	7380
tccatttctt	catcttcact	ctcatcatca	tcaccaccac	catgatgatt	ctcctcctct	7440
tcacgtaacc	tagcaatctc	actctgagct	ctatcaacaa	tetgettett	ctgcaactcc	7500
aaatctctct	gaaaatcagc	teteatette	tccaactcct	tcatttgcto	tttcttactc	7560
ttctccatct	tctcataaac	cttcccaaac	ctctcaacag	aatccgccaa	catcttatac	7620
gaagcagcgt	cattaacctt	cttcctctcg	tactcaacct	catcatcctc	atcctcctcc	7680
tcttcagaat	caccaggact	atccatcatc	tcatcaaaco	cattagactt	atctaaataa	7740
accttagtgt	tcataaacac	: aaactcacct	gaatcaacac	cacaagctaa	a acctaaatcc	7800
gacttgggcg	, aaacacaaag	caacatatco	c aacttattga	a aaaacgacca	a tttacttgaa	7860
cctaaacctg	, atttctcaac	cttaatctt	c tettttetat	acttcctctt	caagtcatca	7920
atcattctcc	tacattgcgt	ctcagattt	c tocatcotta	a gctcctcact	cactttctca	7980
gctacttcat	tccaatcct	gttcctcaaa	a ctccttcta	c ccaattgcaa	a aaacctatct	8040
ccccaaactt	caagcaaca	c aa	,			8062

	•	•		
,	•		,	
,				
	•			
į				
				,
				•
				•
•				
	·			
-				